



## X-ray properties of the “Composite” Seyfert/Star-forming galaxies

A. Wolter<sup>1</sup>, G. Trinchieri<sup>1</sup>, R. Della Ceca<sup>1</sup>, F. Panessa<sup>2,3</sup>, M. Dadina<sup>2</sup>, L. Bassani<sup>2</sup>,  
M. Cappi<sup>2</sup>, A. Fruscione<sup>3</sup>, S. Pellegrini<sup>4</sup>, and G. Palumbo<sup>4</sup>

<sup>1</sup> Osservatorio Astronomico di Brera via Brera 28, 20121 MILANO e-mail:  
anna@brera.mi.astro.it

<sup>2</sup> INAF/IASF, via P. Gobetti 101, 40129 Bologna

<sup>3</sup> Center for Astrophysics, 60 Garden St., Cambridge, MASS

<sup>4</sup> Dipartimento di Astronomia, Università di Bologna, via Ranzani 1, 40127 Bologna

**Abstract.** An enigmatic, small class of IR and X-ray luminous sources, named “Composite” starburst/Seyfert galaxies, has been defined from IRAS and RASS data. The objects have optical spectra dominated by the features of HII galaxies (plus, in some cases, weak Seyfert signatures) but X-ray luminosities higher than expected from starbursts and more typical of Seyfert nuclei. The true nature of this class of objects is still unknown. We present Chandra data of four of these galaxies that were obtained to investigate the nature of the X-ray source. The X-ray spectrum, the lack of any significant extended component, and the observed variability indicate that the AGN is the dominant component in the X-ray domain.

**Key words.** Galaxies: active – Galaxies: starburst – X-rays: galaxies

### 1. Introduction

An enigmatic class of six low redshift galaxies has been defined during a spectroscopic survey of bright IRAS and RASS sources (Moran et al. 1996). They were named “Composite” to indicate their dual nature: the optical spectrum typical of a star forming galaxy is characterized by additional weak broad  $H\alpha$  wings that, together with the high X-ray luminosities, suggests the presence of an AGN. We have performed a study with Chandra of the four “Composites” that still lack a high resolution X-ray observation, with two main goals:


*Send offprint requests to:* A. Wolter, OAB

*Correspondence to:* via Brera, 28, 20121 MILANO

1) to investigate any extended X-ray component possibly associated with the InfraRed Starburst; 2) to determine the nature of the nuclear emission. We present here the first results obtained, while a detailed description of the X-ray data will appear in Panessa et al. (in preparation).

### 2. X-ray data

The sources were all observed with the ACIS-I instrument on board Chandra for a nominal exposure of  $\sim 25$  ksec. Relevant data for the sources are listed in Table 1. Based on the RASS fluxes we had requested a Chandra configuration that would allow more frequent




**Fig. 1.** The light curve of IRAS20069+5929, in bins of 1000 sec, and residuals with respect to a constant (in  $\sigma$ ); the fit  $\chi^2$  is 46.8/25, indicating that variability is present.

read-out of the data by using a smaller field of view to avoid pile-up. The results however indicate that the precautions taken were not necessary, since the observed count rates (see Table 1) are well below the expected value and the pile-up is not significant.

### 2.1. Variability

We compare the total Chandra luminosities to previous RASS data (Moran et al. 1996). We find consistency only for IRAS20069+5929. The luminosity is smaller by a factor  $\sim 10$  for IRAS04392-0123, and a factor  $\sim 2$  for IRAS01072+4954 & IRAS01319-1604, as indicated by R in Table 1. A discrepancy of a factor  $\sim 25$  was also reported for IRAS00317-2142 (Georgantopoulos et al. 2003), while no information is available so far for IRAS20051-1117 that has been observed by Chandra and XMM-Newton. The long term variability alone already indicates that the dominant source of emission is due to a point sources.

We have further investigated short term variability. We compute light curves by extracting the source counts in a region of  $4''$  radius and background in a larger, source free, annulus around it. We bin light curves at



**Fig. 2.** Profile of IRAS20069+5929 (squares) compared to the *chart*-derived PSF with the same spectral distribution of the source (histogram). X-axis is in arcsec.

1000 sec to have enough statistics per bin. We show in Figure 1 the non constant light curve of source IRAS20069+5929, which has the largest statistics of the 4 Chandra observations. We highlight this variability detection since it indicates the presence of a point source even if the average luminosity of this source has not changed since RASS.

### 2.2. Spatial distribution

No extent is evident in any of the four sources, although a low surface brightness component might be present; more detailed analysis is underway. With the *ciao* task *chart* we simulate a point source distribution of photons that has the same spectrum of the observed source. We then extract the profile of both the source the Chandra PSF to compare them. We plot the histograms in Figure 2 for the brightest source IRAS20069+5929 and verify that it is consistent with a point source.


### 2.3. X-ray spectra

We extract spectra for the four sources and fit them with a simple model: a power law modified by low energy absorption due to our galaxy (from Dickey & Lockman 1990) plus an intrinsic redshifted component such as that ex-

**Table 1.** Properties of the 4 “Composite” galaxies under study.

Name	Redshift	c/s ACIS-I	R*	$L_X(2-10\text{keV})$ cgs (Chandra)	L(FIR) cgs
IRAS01072+4954	0.0237	0.063±0.002	0.5	$3.85 \times 10^{41}$	$1.22 \times 10^{44}$
IRAS01319-1604	0.0199	0.115±0.002	0.5	$6.99 \times 10^{41}$	$1.97 \times 10^{44}$
IRAS04392-0123	0.0289	0.015±0.001	0.1	$3.68 \times 10^{41}$	$1.81 \times 10^{44}$
IRAS20069+5929	0.0374	0.180±0.003	1.0	$6.84 \times 10^{42}$	$7.52 \times 10^{44}$

\* R is the ratio between Chandra and RASS fluxes.



**Fig. 3.** The X-ray hard luminosity from Chandra vs. the FIR luminosity from Moran et al. (1996) (circles). The cross represents IRAS 20005-1117 (from Moran et al. 1996, X-ray from RASS); circle+cross is IRAS 00317-2142 (from Georgantopoulos et al. 2003, X-ray from Chandra). Dotted and dashed lines from Persic et al. 2004, solid line from David et al. 1992. See text for details.

pected in the hypothesis that the bright broad component in the optical spectrum is not seen because of absorption. We find instead that intrinsic  $N_H$  is always small or negligible: IRAS01072+4954 and IRAS01319-1604 do not require statistically an intrinsic  $N_H$  component; IRAS04392-0123 has  $N_H^{\text{intrinsic}} \leq 1.17 \times 10^{21} \text{ cm}^{-2}$ ; IRAS20069+5929 has  $N_H^{\text{intrinsic}} = 2.69[2.21 - 3.22] \times 10^{21} \text{ cm}^{-2}$ .

A similar behavior is reported for IRAS 00317-2142, with ASCA data (Georgantopoulos 2000), where there is

no evidence of intrinsic absorption above a few  $10^{20} \text{ cm}^{-2}$ .

In spite of the lower luminosities in the most recent measurements, the current values are still not consistent with the hypothesis of them being produced by the starburst seen at InfraRed wavelengths. This is evident in Figure 3, where we plot (circles) the X-ray hard luminosity from Chandra vs. the FIR luminosity from Moran et al. (1996). The cross represents IRAS 20005-1117 (from Moran et al. 1996, X-rays from RASS); circle+cross is IRAS 00317-2142 (from Georgantopoulos et al. 2003, X-rays from Chandra). From the FIR Luminosity we compute the expected Star Formation Rate ( $\text{SFR} = 8\text{-}80 \text{ M}_\odot/\text{yr}$ ). From the SFR we derive the expected X-ray luminosity according to Persic et al. (2004) (dotted line is contribution from HMXB; dashed line is the total from the Starburst) or David et al. (1992) (solid line). The currently observed X-ray luminosities of the sources are still at least about a factor of 10 higher than both estimates. The AGN dominates in X-rays or the starburst component is anomalous.

### 3. Conclusions

From the X-ray Chandra observations we conclude that the sources are: 1) point-like in the Chandra images, although we cannot exclude a low surface brightness component; 2) variable - both within the single Chandra observation (as in the case of IRAS20069+5929) and with respect to older data; 3) a thermal component is in general not present, even if in a few cases it might account for part of the soft emission; 4) the intrinsic absorption measured in the X-ray band is small or not required by the fit; 5) the current  $L_X$  is still at least an order of magnitude

higher than the predictions from the starburst power, which is of the order of  $\text{SFR} = 10\text{-}100 M_{\odot}/\text{yr}$ .

Since the emission lines from the AGN are weak in the optical band, and absorption is not detected in X-rays, we cannot suggest obscuration as responsible for hiding the optical AGN.

Therefore we conclude that an AGN is present and dominant in X rays, but probably of low luminosity:  $L_X(2\text{-}10 \text{ keV}) = 3.5 \times 10^{41} - 7 \times 10^{42} \text{ erg/s}$ . This could explain the optical faintness observed by Moran et al. (1996).

*Acknowledgements.* We acknowledge partial financial support from Agenzia Spaziale Italiana (ASI).

## References

- David L.P., Jones C., & Forman W., 1992, ApJ, 388, 82  
Dickey J.M. & Lockman F.J., 1990, ARAA, 28, 215  
Georgantopoulos, I., 2000, MNRAS, 315, 77  
Georgantopoulos, I., Zezas, A., Ward, M. J., 2003, ApJ, 584, 129  
Moran, E., Halpern, Jules P., Helfand, David J., 1996, ApJS, 106, 341  
Persic, M., Rephaeli, Y., Braitto, V., Cappi, M., Della Ceca, R., Franceschini, A., Gruber, D. E., 2004, A&A 419, 849