



# SYNTHE spectral calculations matched to high-resolution UV/optical observations

R. Peterson<sup>1,2</sup>

<sup>1</sup> UCO/Lick Observatory, Department of Astronomy, University of California, Santa Cruz, CA 95064 e-mail: [peterston@ucolick.org](mailto:peterston@ucolick.org)

<sup>2</sup> Astrophysical Advances, Palo Alto, CA 944301

**Abstract.** Progress is shown in matching SYNTHE spectral calculations to the observed high-resolution spectra of standard stars. In the mid-UV these are the Sun, Procyon, and solar-type stars ranging in metallicity. Redward of 3000Å, the K giant Arcturus is added. In the mid-UV, from 2300Å to 3000Å, good reproduction of individual spectral absorption lines in solar-type stars is achieved to an iron abundance  $[Fe/H] = -0.5$ . At higher metallicities, many atomic lines appear which are unidentified in the laboratory and so are missing from the calculations. We are fixing this with STIS echelle spectra already in hand. In the near-UV, 3000Å–3800Å, good agreement is found at solar metallicity for solar-type stars, but missing atomic lines pose problems for Arcturus. In the optical, the fit to Arcturus is not good from 4090Å to about 4500Å, due to lines of the molecule SiH, whose wavelengths are poorly known. Beyond 4500Å, excellent agreement is seen in regions of low to moderate telluric molecular absorption, for all stars considered.

**Key words.** Stars: abundances – Stars: atmospheres – Stars: abundances

## 1. Background

My NASA Long-Term Space Astrophysics and Hubble Treasury programs, now in their third year, aim to provide theoretical templates for interpreting the age, metallicity, and populations of galaxies and stellar clusters as young as 1 Gyr from their ultraviolet (UV) and optical spectra. The first task is to calculate stellar spectra as precisely as possible. The first two years were designed to improve the list of line parameters input to spectral calculations, to the point where the spectra of stars of F type and later can be reproduced reliably line-by-line. This paper presents the

status of this effort. This work will provide both theoretical and observations products to the community through the Space Telescope Science Institute's MAST archive service at <http://archive.stsci.edu>. The theoretical products will include extensive grids of spectral calculations, as well as high-resolution atlases comparing calculations to standard stars. Shown here are several atlas comparisons.

## 2. Observational Data

To improve the spectral calculations and to ensure their validity, we match calculations to real spectra of standard stars. Improvements are best done using spectra of high resolution and high signal-to-noise covering as wide

---

*Send offprint requests to:* R. Peterson



didates from the lab line list. We repeat all calculations and comparisons, iterating until a satisfactory match is achieved with the same set of gf values for all stars simultaneously. We assume local thermal equilibrium (LTE), and adopt the ODFNEW models of Castelli & Kurucz (2003). We first redetermine the standards' parameters by demanding matches for all stars in the profiles of strong lines and H $\alpha$  as well as in the weak-line strengths and in mid-UV flux levels. Our Arcturus and Procyon values determined in this way agree well with recent results. The missing lines we add as Fe I, or Ti I if the feature is stronger in HD 184499 than HD 157466. Their line strengths are similar, but HD 184499 has the typical halo  $\alpha$ -element enhancement, while the disk star HD 157466 has scaled-solar abundances.

#### 4. Plots of atlases

Plots are shown comparing our theoretical spectra (light black line) with observational echelle spectra (heavy black line). In color appear theoretical calculations for a single element only. Wavelength in Å appears at the bottom, and Y-axis ticks represent 10% of the full scale. Stars are offset for visibility. Identifications for the strongest lines appear at the top. The values are (1) the decimal digits of the wavelength in Å, (2) a colon if it is a missing Fe I line, (3) the species and its excitation potential in electron volts (or band), and (4) the residual intensity in the solar or Arcturus spectrum. Next to each star's name are best-fit model parameters  $T_{\text{eff}}$ ,  $\log g$ , [Fe/H] (log of the stellar-to-solar iron abundance ratio), and microturbulent velocity in  $\text{km s}^{-1}$ . In the mid-UV, a single flux normalization constant is adopted at all wavelengths. This is possible because the STIS spectra are fluxed. It provides a critical constraint on the modeling. We find a normalization constant for Procyon

that is 8.5% larger than that implied by its angular diameter, marginally consistent with the current accuracy of STIS spectroscopic fluxes.

*Acknowledgements.* We thank Dr. Kurucz and Dr. Castelli for making their atlases, line lists and model grids available. By the kind permission of the Astronomical Society of the Pacific Conference Series, we plot the Arcturus data from Hinkle et al. (2000). We thank the STScI TAC for its allocation of observing time to the Treasury program. We thank the UVES Paranal Observatory Project for making its spectra of Arcturus available. We are indebted to Jon Fulbright, Debra Fischer, Steve Vogt, Mike Bolte, and Jennifer Johnson for generously contributing their spectra towards this end. This work has been supported by grants from the Space Telescope Science Institute (GO 9455/9974/10094) and the NASA Long-Term Space Astrophysics program.

#### References

- Bagnulo, S., Jehin, E., Ledoux, C., Cabanac, R., Melo, C., Gilmozzi, R., & The ESO Paranal Science Operations Team 2003, *The Messenger*, 114, 10, <http://www.sc.eso.org/santiago/uvespop/>
- Castelli, F., & Kurucz, R. L., 2003, *Proceed. of IAU Symp. 210, Modelling of Stellar Atmospheres*, eds. N. Piskunov et al., poster A20 on the enclosed CD-ROM, (astro-ph/0405087)
- Hinkle, K., Wallace, L., Valenti, J., & Harmer, D. 2000, *Visible and Near Infrared Atlas of the Arcturus Spectrum 3727-9300 Å* ed. Kenneth Hinkle, Lloyd Wallace, Jeff Valenti, and Dianne Harmer. (San Francisco: ASP) ISBN: 1-58381-037-4, 2000.,
- Kurucz, R. L., Furenlid, I., & Brault, J. T. L. 1984, *National Solar Observatory Atlas, Sunspot*, New Mexico: National Solar Observatory, 1984,
- Peterson, R. C., Dorman, B., & Rood, R. T. 2001, *ApJ*, 559, 372