

# Abundance analysis of the chemically peculiar star HD 74169 (A0IVp)

E. Niemczura and G. Połubek

Astronomical Institute, Wrocław University, Kopernika 11, Wrocław, Poland e-mail: eniem@astro.uni.wroc.pl

**Abstract.** We present the results of the abundance analysis of the chemically peculiar star HD 74169 based on the echelle spectrum obtained with the spectrograph UVES. This high resolution and high signal-to-noise observations are compared with a synthetic spectrum computed with the SYNTHE code. As an input atmosphere model for SYNTHE we used an ATLAS 9 model with effective temperature  $T_{\text{eff}} = 10000$  K and surface gravity  $\log g = 4.0$  dex. The iterative procedure used to derive stellar individual abundances allows us to take into account even blended lines. The best fit to the observed spectral line profiles was obtained with a rotational velocity  $V_e \sin i = 20$  km s<sup>-1</sup> and turbulence  $V_{\text{turb}} = 2.1$  km s<sup>-1</sup>. The average abundances of HD 74169 are typical for an Ap star of Cr type: C is deficient, Cr and Fe are strongly overabundant. The analyzed heavy and rare-earth elements are also overabundant.

**Key words.** Stars: abundances – Stars: atmospheres

## 1. Introduction

HD 74169 (KR Vel, A0IVp) is one of the chemically peculiar stars in the field of the open cluster IC 2391 (*o* Vel). The first information about peculiarity of this star appeared in the sixties (Bertaud 1965). Buscombe (1965) and Levato and Malaroda (1984) classified HD 74169 as Ap EuCr(Sr). The KR Vel membership in the young cluster IC 2391 was confirmed by Pöhlh et al. (2003). They analyzed the evolutionary status of CP2 stars of this cluster and found that both the location of HD 74169 on the H-R diagram and its age are within the expected error of the overall age of the cluster. Also Dodd (2004) confirmed the cluster membership of this star on the ba-

sis of proper motion components as well as infrared multicolour photometry. Furthermore, HD 74169 was suggested by North (1984) as a variable star with the period equal to 4.<sup>d</sup>59 obtained on the basis of lightcurves determined in the Geneva photometric system.


In this paper we present the first attempt of the abundance analysis of this star. High quality observations are described in Sect. 2. The iterative method of the abundance analysis and determination of effective temperature and surface gravity are presented in Sect. 3. Results and conclusions are given in Sect. 4.

## 2. Observations

The spectrum of HD 74169 was obtained on February 8, 2001 at ESO with the VLT UV-Visual Echelle Spectrograph UVES at Unit

---

Send offprint requests to: E. Niemczura



**Fig. 1.** Numerical derivatives used by the spectral synthesis method  $\partial_x F_\lambda$ , where  $x$  denotes various parameters involved in stellar spectrum.

2 Kueyen. The slit width was set to 0.5'' which provides a spectral resolution about 80,000. The data were observed and reduced by Bagnulo et al. (2003) and are part of *the Library of High-Resolution Spectra of stars across the H-R Diagram*. The analyzed spectrum covers the spectral range from about 3,000 to 10,000 Å.

### 3. Spectrum Synthesis Method


The synthetic spectra were computed with the SYNTH code (Kurucz 1993a) ported under GNU Linux by Sbordone et al. (2005). All the atmospheric models were computed with the line-blanketed LTE ATLAS9 code, which handles line opacity with the opacity distribution function method (ODF). Also the stellar lines were identified on the basis of the Kurucz lines list (Kurucz 1993b).

For the analysis of the data an efficient spectral synthesis method based on the least-squares optimization algorithm was used (Takeda 1995, Bevington 1969). This method allows for the simultaneous determination of various parameters involved with stellar spectra and consists in minimization of the deviation between theoretical flux distribution and

the observed one. The synthetic spectrum depends on the stellar parameters like effective temperature  $T_{\text{eff}}$ , surface gravity  $\log g$ , rotational velocity  $V_e \sin i$ , turbulence  $V_{\text{turb}}$ , radial velocity  $V_r$  and relative abundances of the elements  $\epsilon_i$ , where  $i$  denotes individual element. The first two parameters were not determined during the iteration process but were considered as the input ones. All the other above-mentioned parameters can be determined simultaneously because they produce detectable and different spectral signatures (see Fig. 1). The theoretical spectrum was fitted to the normalized observed one. The continuum was subjectively drawn by connecting the highest points of the analyzed spectrum part. As an example of the spectral synthesis method we show a comparison of theoretical and observed spectrum for first six iteration steps in the spectral range 3602 – 3607 Å (Fig. 2).

#### 3.1. Input parameters $T_{\text{eff}}$ and $\log g$

The determination of the effective temperature for Ap stars is a complicated problem. The effective temperature  $T_{\text{eff}} = 9950$  K of HD 74169 has been previously obtained by Rode-Paunzen et al. (2004) from Geneva pho-



**Fig. 2.** Part of the observed flux together with synthetic one for six iteration steps is plotted. In this spectral range the abundances of iron, chromium and nitrogen as well as radial velocity were determined.

tometry. Another estimation of the effective temperature was done by Glagolevskij (1994) who used the Shallis-Blackwell (Blackwell et al. 1980) method starting with the total flux of the energy emitted by the star and found the average temperature 9650 K. Glagolevskij gave also two other values,  $T_{\text{eff}} = 9700$  K derived from reddening free index  $Q$  (Johnson and Morgan 1953) and  $T_{\text{eff}} = 10100$  K from  $X$  parameter. Applying the Napivotzki et al. (1993) calibration to the observed  $uvby\beta$  photometry extracted from the WEBDA database (Mermilliod 2005) we obtained two values of  $T_{\text{eff}}$  from indices  $[c_1]$  ( $T_{\text{eff}} = 10560$  K) and  $[u-b]$  ( $T_{\text{eff}} = 10050$  K). We also used Hauck and North (1993) calibration of the Geneva photometry to derive  $T_{\text{eff}} = 9620$  K from  $(B2 - G)$  index. As one can see there is a significant discrepancy between all these values of effective

temperature. In our analysis we adopted  $T_{\text{eff}} = 10,000$  K as close to the mean value obtained on the basis of all mentioned methods.


The logarithmic surface gravity was determined using theoretical profile fitting to the hydrogen Balmer lines,  $H_\beta$ ,  $H_\gamma$  and  $H_\delta$ . The comparison between the observed and calculated  $H_\gamma$  profiles is presented in Fig. 3. We were searching for the best fit for the temperatures from 9500 to 10500 K with the step equal to 50 K. Considering the problem in the continuum normalization of the hydrogen lines we adopted surface gravity of 4.0 dex as the appropriate value for this star. The profile is better fitted for lower temperatures, what may be taken as evidence in favour of the value obtained by us. Errors in adopted parameters are estimated to be 200 K for the effective temperature and 0.2 dex in  $\log g$ .

#### 4. Results and conclusions


About 300 lines were analyzed in the spectrum of the chemically peculiar star HD 74169. The majority of them were identified as iron group elements features. We were able to determine abundances of 26 elements. Only a few of them have lower abundances than the Sun. The results presented here provide the mean abundances without a separate analysis of the neutral and ionized species. This is caused by the fact that in many cases the separation of the influence of neutral and ionized state of the same element to the line profile is not possible. The mean abundances derived for the atmosphere of HD 74169 are given in Table 1. In the same table the solar values from Grevesse and Sauval (1998) are shown for comparison.

The rotational velocity was determined as  $\approx 20$  km s $^{-1}$  from the analysis of many iron group elements lines. From the scatter of the values determined for different lines we can assume the error of this parameter as 2 km s $^{-1}$ . The obtained value of  $V_e \sin i$  is lower than the 40 km s $^{-1}$  given by Levato and Garcia (1984).

We estimate the mean turbulence velocity  $V_{\text{turb}}$  as equal to 2.1 km s $^{-1}$  by fitting simultaneously the element abundances and  $V_{\text{turb}}$  for some strong lines.



**Fig. 3.** The observed  $H_{\gamma}$  profile for HD 74169 is plotted with theoretical LTE profile obtained for the atmospheric model with  $T_{\text{eff}} = 10000$  K,  $\log g = 4.0$  dex and the chemical abundances determined in this paper.



**Fig. 4.** Abundances of HD 74169 for light elements, iron group elements, heavy and rare earth elements, compared to the results for other chemically peculiar stars (see text).

#### 4.1. Light elements

Lines of C, N, Ne, Na, Mg, Si, P, S and Ar were measured and their abundances were

computed. Only the abundance of carbon is lower than the solar value. The abundances of N, Ne, Na, Mg, Si, P, S and Ar are overabundant. All these values were determined on the

**Table 1.** Atmospheric abundances (third column)  $\log \varepsilon$  for HD 74169. In the second column the number of features used to the chemical analysis is presented while in the last one the solar abundances from Grevesse and Sauval (1998) are showed.

Element	n	$\log \varepsilon$	$\log \varepsilon_{\odot}$
C	4	-3.77	-3.52
N	3	-3.51	-4.12
Ne	1	-2.60	-3.96
Na	2	-3.20	-5.71
Mg	6	-4.10	-4.46
Si	12	-4.22	-4.49
P	1	-5.72	-6.59
S	4	-4.47	-4.71
Ar	3	-3.02	-5.48
Ca	8	-4.91	-5.68
Sc	8	-9.12	-8.87
Ti	28	-6.50	-7.02
V	19	-7.34	-8.04
Cr	77	-4.31	-6.37
Mn	26	-5.51	-6.65
Fe	135	-3.60	-4.54
Co	5	-6.02	-7.12
Ni	26	-5.41	-5.79
Cu	1	-6.59	-7.83
Zn	1	-6.27	-7.44
Y	2	-8.42	-9.80
Zr	2	-8.40	-9.44
Ba	1	-10.18	-9.91
Ce	4	-7.83	-10.46
Nd	1	-7.20	-10.54

basis of weak features blending with strong iron peak elements. Only for magnesium and silicon more than four lines were detected and in these cases the mean values are better determined. The other values should be treated with caution. The standard deviation  $\sigma$  for magnesium and silicon abundances are 0.55 dex and 0.85 dex respectively. Large values of standard deviations result from the fact that only mean abundances are derived without a separate assessment of the neutral and ionized species.

#### 4.2. Iron group elements

The spectrum is dominated by the iron peak elements. Except for Co, Cu and Zn the abundances are determined from more than five features. Only Sc (-9.10 dex) has a lower abundance in comparison to the Sun. All other elements are found to be overabundant by more than 0.4 dex. Iron shows a high abundance of -3.60 dex which is significantly greater than the solar value. Another remarkable result is the more than 2.0 dex overabundance of chromium. The standard deviations are relatively small (about 0.35 dex) for Cr, Mn, Fe and Co. For the first three elements this is the effect of many analyzed lines. The existing discrepancies for these and other lines might be explained after the introduction of vertical stratifications for the elements. Higher values of  $\sigma$  for Ca, Ti, V, Ni and Sc first of all can result from the fact that the mean values are computed from both neutral and ionized species. Additionally in the case of Sc and V only the weak lines or blends were analyzed.

#### 4.3. Heavy and rare earth elements

Because of the relatively high rotational velocity of the analyzed star, very few signatures of the heavy and rare earth elements were found as a separate line or blend in the spectrum. In most cases only the influence of such elements to the wings of strong lines were seen. The method we use let us determine the abundance even in these cases. We were able to determine abundances of Y, Zr, Ba as well as rare earth elements: Ce and Nd, identified only on the basis of the Kurucz lines list. The abundances of yttrium (-8.42 dex) and zirconium (-8.40 dex) are enhanced by more than 1.0 dex in comparison with the solar values, whereas barium (-10.18 dex) has a lower abundance. However, the barium abundance was determined on the basis of only one weak line which blends with a relatively strong Fe line. The rare earth elements, cerium (-7.83) and neodymium (-7.20) show overabundances by more than 2.0 dex. HD 74169 seems to be very enhanced in these elements in comparison to the Sun. The same conclusions were deduced by Kato (2003) for

another silicon star HD 170973. The list of identified heavy and rare earth elements would be more complete if other sources such as VALD (Kupka et al. 1999) or the DREAM (Biemont et al. 1999) databases were included in the analysis.

#### 4.4. Errors in the abundance analysis

Derived abundances contain errors connected with a number of sources of uncertainty, including the adopted atomic data, most importantly the oscillator strengths, the adopted atmospheric parameters and model, and the observed spectra. For some sources we can directly estimate the associated errors. The abundance differences due to the change of  $T_{\text{eff}}$  by 200 K show the discrepancy of about 0.25 dex, whereas the change of  $\log g$  by 0.2 dex cause the differences within 0.1 dex. The values of the differences depend on the analyzed element.

#### 4.5. Comparison with other stars

Figure 4 shows the comparison of the abundances of light, iron peak, heavy and rare earth elements obtained for HD 74169, with the literature values for the Sun (Grevesse and Sauval 1998), HD 48915, HD 170672, HD 43819, HD 133029, HD 192913 (Saffe and Levato 2004), HD 116458 (Nishimura et al. 2004) and HD 170973 (Kato 2003). Except for HD 48915 (Sirius) and HD 170672 (Vega) all stars are chemically peculiar silicon stars. The abundance pattern of HD 74169 is consistent with the results for other stars, especially for iron peak elements. The abundances of light elements Ne and Ar were determined only for HD 74169.

In this paper we present preliminary results. In the next step we are going to show a more detailed abundance determinations with a separate analysis of the neutral and ionized elements for all normal and chemically peculiar stars of IC 2391.

*Acknowledgements.* This work was partially supported by the KBN grant No. 1/P03D/01627.

## References

- Bagnulo, S., Jehin, E., Ledoux, C., et al. 2003, *The Messenger*, 10
- Bertaud, C. 1965, *JO*, 48, 211
- Bevington, P.R. 1969, in *Data Reduction and Error Analysis for the Physical Science* (New York: McGraw-Hill)
- Biemont, E., Palmeri, P., Quinet, P. 1999, *Astrophys. Space Sci.*, 635, 269
- Blackwell, D.E., Petford, A.D., Shallis, M.J. 1980, *A&A*, 82, 249
- Buscombe, W. 1965, *MNRAS*, 129, 411
- Dodd, R.J. 2004, *MNRAS*, 355, 959
- Glagolevskij, Yu.V. 1994, *BSAO*, 38, 152
- Grevesse, N., Sauval, A.J. 1998, *Space Sci. Rev.*, 85, 161
- Hauck, B., North, P. 1993, *A&A*, 269, 403
- Johnson, H.L., Morgan, W.W. 1953, *ApJ*, 117, 313
- Kato, K. 2003, *PASJ*, 55, 1133
- Kupka, F., Piskunov, N., Ryabchikova, T.A. 1999, *A&AS*, 138, 119
- Kurucz, R., 1993a, CD-ROM, No. 18, <http://kurucz.harvard.edu>
- Kurucz, R., 1993b, CD-ROM, No. 13, <http://kurucz.harvard.edu>
- Levato, H., Garcia, B. 1984, *ApL*, 24, 49
- Levato, H., Malaroda, S. 1984, *ApL*, 24, 37
- Mermilliod, J.-C. 2005, <http://obswww.unige.ch/webda/>
- Napiwotzki, R., Schönberner, D., Wenske, V. 1993, *A&A*, 268, 653
- Nishimura, M., Sadakane, K., Kato, K., et al. 2004, *A&A*, 420, 673
- North, P. 1984, *A&AS*, 55, 259
- Pöhl, H., Maitzen, H.M., Paunzen, E. 2003, *A&A*, 402, 247
- Rode-Paunzen, M., Pöhl, H., Paunzen, E., Maitzen, H.M. 2004, *IAU Symp. No. 224*
- Saffe, C., Levato, H. 2004, *A&A*, 418, 1083
- Sbordone, L., Bonifacio, P., Castelli, F. 2005, <http://wwwuser.oat.ts.astro.it/atmos/>
- Takeda, Y. 1995, *PASJ*, 47, 287