

Properties of the base of streamers from UV and EUV observations

S.Parenti¹, E. Landi², B.J.I. Bromage³

¹ Dipt. di Astronomia e Scienza dello Spazio, Università di Firenze, Largo E. Fermi 3, 50125 Firenze

² Artep, Inc. Columbia MD 21044 at Naval Research Laboratory, Washington DC 20375-5320, USA

³ Centre for Astrophysics, University of Central Lancashire, Preston, UK

Abstract.

In this paper we investigate the physical parameters of a streamer observed during the Spring 2000 quadrature of SOHO, the Sun and Ulysses. The analysis was carried out using SOHO/CDS and SOHO/SUMER coordinated observations of the low corona. We analysed six days of observations of the same region, deriving electron temperature, density, emission measure and composition. We found full agreement in the results from the two instruments. We conclude that we were observing an omogeneous streamer, with coronal composition and not completely isothermal.

Key words. Sun – EUV diagnostic

1. Introduction

The configuration of quadrature reached twice times per year by the SOLar and Heliospheric Observatory (SOHO) and Ulysses satellites gives the oppurtunity of make coordinated observations to investigate the solar plasma during its flow from the corona to the interplanetary medium (Suess & Poletto 2001). This kind of observations are very important because they allow to investigate the variation of the physical parameters with the solar distance, giving information on the origin, on the Sun, of the solar wind.

Send offprint requests to: S. Parenti

Correspondence to: Largo E. Fermi 3, 50125 Firenze, IT

The understanding of the variation of the elements abundance in the solar athmospere and solar wind, are at the base of all unsolved problems in solar physics. Several studies have been carried out on this topic (i.e. Meyer 1996, Fludra et al. 1999, Raymond et al. 2001), but the problem is still opened. In addition the previous studies on the composition in corona, we mention the depletion found in streamers at $1.5 R_{\odot}$ with respect to the photospheric values (Raymond et al. 1997, Parenti et al. 2000), which was not visible in the low corona (Feldman et al. 1998). The electron temperature measured in streamers show that they are often isothermal (Raymond et al. 1997, Feldman et al. 1998). The density in these

structures has been found to decrease with the solar distance according to the hydrostatic law (Gibson et al. 1999). We present a summary of the results for the radial and temporal variation of electron temperature, density and composition in a streamer, obtained with the SUMER (Wilhelm et al. 1995) and CDS (Harrison et al. 1995) instruments on board of SOHO. In particular we concentrate the study along the Ulysses direction. The full description of the present analysis can be found in Parenti et al. (2003). Results from other instruments which participate to the quadrature observation can be found in Bemporad et al. (2003)

2. The Observations

The observations were carried out between the 13 and 18 June 2000, pointing the two instruments on the radial to Ulysses at -58.2° in the South-East quadrant. CDS observations consisted of full NIS spectra, obtained rastering the corona with the $4'' \times 120''$ slit up to $1.3 R_\odot$. The SUMER observations were taken pointing the slit number 1 at the Ulysses radial at $1.2 R_\odot$. A full detector A spectra was obtained for each spatial position. Figure 1 shows the solar corona as seen by the EIT 195 Å channel on June 17 with the CDS rasters and the SUMER slit superimposed. Full details of the observation can be found in Parenti et al. 2003.

A careful inspection of the solar corona during the time of observations has shown the Ulysses direction to cross a streamer which seems to be present for all the days of the observation.

3. The Results

The following analysis was carried out applying the spectroscopic diagnostic techniques for optically thin lines (Mason & Monsignori Fossi 1994). We used the the CHIANTI database (Dere et al. 1997, Young et al. 2003),




Fig. 1. SOHO/EIT images at 195 Å with the SUMER slit and the CDS/NIS rasters superimposed, for the June 13 (left) and 17 (right). The rasters are represented by the white rectangles. The SUMER slit is shown as a thick white segment. The arrow indicates the Ulysses direction at a latitude -58.2° .

assuming Mazzotta et al. (1998) ionization equilibrium and adopting the coronal composition of Feldman & Laming (2000).

Selected areas of the CDS rasters were used to derive a radial profile of electron temperature and density along the Ulysses direction. Figure 2 shows an example of the results for temperature. These profiles were obtained applying the line ratio technique. The CDS data show the streamer to have a double component in temperature. This result was also confirmed by an Emission Measure (EM) analysis, which suggested a two temperature plasma, one at about 1.4×10^6 K and 1.9×10^6 K at $1.2 R_\odot$. A similar analysis was carried out on the SUMER data. However, due to the slit position, CDS and SUMER field of view intersect only in one point along the Ulysses direction, which corresponds to $1.2 R_\odot$. The EM analysis on SUMER data suggested an isothermal plasma with a temperature almost stable at 1.3×10^6 K over all the days and the latitudes covered by the of




Fig. 2. CDS electron temperature derived from the ratios $[\text{Fe xv}] 360.761/[\text{Fe xiv}] 334.172$ (diamonds), $[\text{Fe xiii}] 348.183/[\text{Fe xii}] 364.467$ (stars) and from $[\text{Mg x}] 609.794/[\text{Mg ix}] 368.070$ (triangles) for 17 June data.

the observations. Both the SUMER and CDS temperature derived at $1.2 R_{\odot}$ (the Ulysses direction) over all the campaign are reported in Figure 3 (Top). We notice a good agreement between the two measurements.

Figure 3 (Middle) reports the electron density derived from the two instruments applying the line ratio technique. The data refer to a distance of $1.2 R_{\odot}$ in the Ulysses direction. The figure shows that the streamer remains homogeneous over all the period of the quadrature. The bottom panel of Figure 3 shows the EM derived over the same period and distance from the two instruments. Also these results show the homogeneity over time of the observed streamer. With the EM analysis we could also study the element composition. Although our data cannot allow to derive the absolute abundances, we could test that the relative elemental abundances are compatible with coronal values for distances up to $1.3 R_{\odot}$.

4. Conclusions

This paper summarises the results we obtained analysing CDS and SUMER data of a streamer in the low corona observed dur-




Fig. 3. SUMER (stars) and CDS (diamonds) temporal variation of density (Top), temperature (Middle) and Emission Measure (Bottom) along the Ulysses direction at $1.20 R_{\odot}$.

ing the Spring 2000 SOHO-Ulysses quadrature. The streamer we were observing showed an almost constant temperature of $1.3 - 1.4 \times 10^6$ K during all the period of our observations. Both the instruments indicate the presence of an isothermal component of the plasma in their field of view which emits at a temperature smaller than 1.6×10^6 K. A second component at about 1.9×10^6 K is seen by CDS. We believe that

this component is not seen in the SUMER data because of the absence of lines emitting at such temperatures, in the waveband covered by the SUMER. Along the Ulysses direction, where the field of view of the two instruments overlap, the measurements of the density are very similar. There is not much change of these quantities during the period of the observation, to indicate that there is no significant change in the structure that is crossed by the fields of view. Differently from SUMER, CDS finds a small increase of EM on June 17 from the contribution of the low temperature plasma. Both CDS and SUMER data indicate a depletion of EM with distance. This result, combined with their density results, along the Ulysses direction and across the streamer, shows that there is a change in the streamer morphology.

Both the instruments can measure the relative abundances of the observed plasma. The results show compatibility with the coronal composition, which indicate that there is an overabundance of the low FIP (First Ionization Potential) elements with respect to photospheric values.

Acknowledgements. EIT images are courtesy of SOHO/EIT consortium. SOHO is a project of international cooperation between ESA and NASA. CHIANTI is a collaborative project involving the NRL (USA), RAL (UK), and the Universities of Florence (Italy) and Cambridge (UK).

References

- Bemporad, A., et al. 2003, ApJ, submitted
 Dere, K.P., et al. 1997, A&AS 125, 149
 Feldman, U., Schühle, U., Widing, K.G., & Laming, J.M. 1998, ApJ 505, 999
 Feldman, U., & Laming, J.M. 2000, Phys. Scr. 61, 222
 Fludra, A., et al. 1999, in "The many faces of the sun: a summary of the results from NASA's Solar Maximum Mission", Springer, p.89
 Gibson, S.E., et al. 1999, Geoph. Res. 104, 9691
 Harrison, R.A., et al. 1995, Sol. Phys. 162, 23
 Mason, H.E., & Monsignori Fossi, B.C. 1994, A&A 6, 123
 Mazzotta, P., Mazzitelli, G., Colafrancesco, S., & Vittorio, N. 1998, A&AS 133, 403
 Meyer, J.P. 1996, "Proceedings of the Second Recontres du Vietnam", J. Tran Thanh Van, et al. editors, October 22-28, 1995. Editions Frontieres, p. 2
 Parenti, S., et al. 2000, A&A 363, 800
 Parenti, S., Landi E., Bromage, B.J.I. 2003, ApJ, in press.
 Raymond, J.C., et al. 1997, Sol. Phys. 175, 645
 Raymond, J.C., et al. 2001, "Joint SOHO/ACE workshop - Solar and Galactic Composition", AIP, 598, 49
 Suess, S., & Poletto, G. 2001, Space Sci. Rev. 97, 59
 Young, P.R., et al. 2003, ApJS 144, 135
 Wilhelm, K., et al. 1995, Sol. Phys. 162, 189