



# XMM-Newton observation of the composite SNR G0.9+0.1 and the discovery of a new transient source

L. Sidoli<sup>1</sup>, F. Bocchino<sup>2</sup>, S. Mereghetti<sup>1</sup> and R. Bandiera<sup>3</sup>

<sup>1</sup> IASF/INAF, via Bassini 15, 20133 Milano, Italy e-mail: sidoli@mi.iasf.cnr.it

<sup>2</sup> INAF-Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, 90134 Palermo, Italy

<sup>3</sup> INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

**Abstract.** We present the preliminary results of a *XMM-Newton* observation of the composite supernova remnant G0.9+0.1 located in the Galactic Center region. The spectrum of the extended pulsar wind nebula can be well fit with a single power-law with a photon index of 1.9 and a large absorbing column density of  $1.4 \times 10^{23} \text{ cm}^{-2}$ . We confirm the X-ray emission previously discovered from the radio shell. We serendipitously discovered a new transient source, likely a new low mass X-ray binary probably located in the Galactic Center region.

**Key words.** SNR – G0.9+0.1

## 1. Introduction

G0.9+0.1 is a composite supernova remnant located in the Galactic Center region. The high interstellar absorption prevented its detection at X-rays (except for a marginal detection with the *Einstein* Observatory, Helfand & Becker 1987) up to the BeppoSAX observation of the region (Mereghetti et al. 1998; Sidoli et al. 1999), where X-ray of non-thermal origin were observed from the radio core of the remnant.

This emission was interpreted as produced by the pulsar wind nebula (PWN) of G0.9+0.1. During a recent *Chandra* observation, the PWN could be spatially resolved. No X-ray pulsations were detected, making the identifi-


cation of the supposed pulsar embedded in the center of the remnant, rather uncertain. Indeed, two possible X-ray sources could host the central neutron star: a faint X-ray source, called CXOU J174722.8-280915, or a clump of X-ray emission coincident with the radio core (Gaensler et al. 2001).

During the first *XMM-Newton* observation of G0.9+0.1 evidence for X-ray emission from the radio shell was found (Porquet et al. 2003). Here we report the preliminary results of the deepest X-ray observation ever performed of this SNR.

## 2. X-ray Emission from the Pulsar Wind Nebula

The entire X-ray emission from the PWN has been studied extracting counts from MOS and

Send offprint requests to: L. Sidoli  
Correspondence to: sidoli@mi.iasf.cnr.it



**Fig. 1.** Spectral softening of the PWN spectrum at larger distances from the core of the nebula

PN cameras. The fit with an absorbed power-law gave  $\Gamma \sim 1.9$ ,  $N_{\text{H}} = 1.4 \times 10^{23} \text{ cm}^{-2}$ , and flux  $F = 4.8 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$  (2–10 keV, corrected for interstellar absorption), translating into a luminosity of  $L_{\text{X}} \sim 5 \times 10^{34} \text{ erg s}^{-1}$  (for a distance of 10 kpc).

A spectral analysis of three different annular regions centered on the PWN peak confirms a softening of the spectrum at larger distance from the peak of the PWN (see Fig. 1).

### 3. X-ray Emission from the Shell

The extraction of counts from a region spatially coincident with the radio shell confirms the presence of net rate from it (after subtracting from a background extracted from the same image and appropriately corrected for the different area and the vignetting).

The fit with an absorbed powerlaw resulted in a steep spectrum with a photon index of  $\sim 3$ , and a flux corrected for the absorption of  $\sim 3 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$  (2–10 keV). Fitting

with a hot plasma model (MEKAL in XSPEC) resulted in a very high temperature of 3 keV (high if compared with the typical temperature of SNR shells).

### 4. A new X-ray Transient

During this *XMM-Newton* observation we serendipitously discovered a new X-ray source (Sidoli & Mereghetti 2003) at the sky position  $R.A. = 17^{\text{h}} 47^{\text{m}} 16.0^{\text{s}}$ ,  $Dec. = -28^{\circ} 10' 45''$  (J2000, 5'' error).

The spectrum of this new source is well fit with an absorbed power-law with photon index  $2.1 \pm 0.1$  and a high column density of  $(8.9 \pm 0.5) \times 10^{22} \text{ cm}^{-2}$ . This suggests that the transient source is located at the GC distance. In this case, its luminosity is about  $5 \times 10^{34} \text{ erg s}^{-1}$  (2–10 keV, corrected for the absorption). The source was not visible in the previous *XMM-Newton* observation performed on September 23, 2000 (Porquet et al. 2003), implying a flux at least a factor 80 fainter.

These properties suggest that this source is a new transient low mass X-ray binary located in the GC region.

### References

- Gaensler B.M., Pivovarov M.J., Garmire G.P. 2001, *ApJ*, 556, L107
- Helfand D.J., Becker R.H., 1987, *ApJ* 314, 203
- Mereghetti S., Sidoli L., Israel G. L. 1998, *A&A*, 331, L77
- Porquet, D., Decourchelle, A., Warwick, R.S. 2003, *A&A*, 401, 197
- Sidoli L., Mereghetti S., Israel G.L., et al. 1999, *ApJ* 525, 215
- Sidoli L., Mereghetti S., 2003, *ATel* n.147