



# Oxygen and Sulfur Enrichment in Low Metallicity HII Galaxies

C. Kehrig<sup>1,2</sup>, J.M. Vílchez<sup>1</sup>, E. Telles<sup>2</sup>, F. Cuisinier<sup>3</sup>, and E. Pérez-Montero<sup>4</sup>

<sup>1</sup> Instituto de Astrofísica de Andalucía – C/ Camino Bajo de Hueter 50, 18008 Granada, Spain e-mail: kehrig@iaa.es

<sup>2</sup> Observatorio Nacional do Rio de Janeiro – Rua General Jose Cristino 77, 224070 Rio de Janeiro, Brasil

<sup>3</sup> GEMAC, Observatório do Valongo/UFRJ Ladeira do Pedro Antônio 43, 20080-090 Rio de Janeiro, Brasil

<sup>4</sup> Universidad Autonoma de Madrid, Cantoblanco, 28049 Madrid, Spain

**Abstract.** We have analysed long-slit spectrophotometric observations of a sample of 35 HII galaxies in the red (6000 Å - 1μm), including the nebular [SIII] lines  $\lambda\lambda 9069, 9532\text{Å}$ . Given the uncertainties associated with the [SIII] $\lambda 6312\text{Å}$  line and the importance of S<sup>++</sup> (the dominant ion) in determining the S/H, it is clear that the observations of the strong near-IR [SIII] lines were needed. All these galaxies were observed previously in the blue wavelength range from which we could derive the O/H abundance using the T<sub>e</sub>[OIII] temperature.

**Key words.** Stars: supernovae – Stars: Population III – Galaxies: low metallicity – Galaxies: abundances

## 1. Introduction

The study of the variation of one chemical element relative to another can provide information on the nucleosynthesis origins of the elements and on the form of the initial mass function (IMF).

Observations of HII dwarf galaxies still provide a direct means of determining the chemical content of the interstellar medium of almost chemically unevolved galaxies, where metal enrichment of the interstellar medium by supernovae has been operating in very low metallicity environments. Sulfur and oxygen are both end-products of explosive nucleosynthesis. The sources of oxygen and its abundance


are well determined. On the other hand, there is little direct observational evidence regarding the most likely stellar mass range for the sulfur production. It should be emphasized also that substantial uncertainties about the sulfur yields still remain.

Thus, the comparison of sulfur and oxygen abundances can provide some clues about the sulfur nucleosynthesis, and the optimum stellar mass range for the production of sulfur.

Another interesting point, somewhat speculative, is related to the possible role played by Population III stars. The study of S/O and, in particular, its possible deviations from the solar ratio at very low metallicities, could be of special interest. It could shed light on the early metal enrichment by Population III stars, several hundred million years after the Big Bang:

---

Send offprint requests to: C. Kehrig



**Fig. 1.** The oxygen and sulfur abundance distribution for our sample of HII galaxies. On the right panel, the dashed and empty histograms show the number of galaxies with S/H derived from Te[SIII] or obtained from  $S_{23}$ , respectively; the oxygen distribution presents only galaxies with O/H derived from Te[OIII].

according to recent zero metallicity stars models the expected value for S/O could reach 1/3 of the solar value (Nomoto et al. 2004).

## 2. Observations

The data base of this work consists of 35 intermediate resolution spectra of HII galaxies in the wavelength range of 6000 Å to 1  $\mu$ m. For six of the 35 observed galaxies it has been possible to calculate the electron temperature  $T_e$ [SIII]. For the galaxies for which we could not measure the auroral [SIII] $\lambda$ 6312Å line, we made use of the empirical abundance indicator  $S_{23}$  (Vilchez & Esteban 1996) to estimate S/H (Pérez-Montero et al. 2005). Besides we have spatially resolved distinct star formation knots and extracted their different spectra for 13 galaxies (Kehrig et al. 2004). All these galaxies were observed previously in the blue wavelength range, from which we could derive the O/H through the  $T_e$ [OIII] (Kehrig 2003).

Figure 1 shows the distribution of oxygen and sulfur abundances for our sample of HII galaxies. We can see that most of the galaxies present abundance values that are below solar. This is the expected behaviour since our sample is composed by low-metallicity galaxies.


## 3. Results and Discussion

A close look to figure 2 reveals that there is a rather large scatter ( $\pm 0.2$  dex) in S/H at a given value of O/H. This may reflect uncertainties due to the correction of the near-IR [SIII] lines from atmospheric absorption or to the faintness of the [SIII] $\lambda$ 6312Å line. Up to 10% of the points present S/O values above  $3\sigma$  of the solar ratio. We are evaluating the contribution of all observational errors to the determination of these points. Nevertheless, at this level of uncertainty, we can conclude that there is no statistical evidence for any systematic variation of S/O with O/H; therefore, our result is in agreement with the current supernovae yields for S and O.


In figure 3, the data show no evidence for any systematic variation of the derived S/H abundance with respect to the ionization structure; confirming the reliability of the empirical method used to derive S/H.

## 4. Conclusions

We present a study of the variation of S/H as a function of O/H in low metallicity environments. The data presented here, together with other studies of S/H based upon the near-IR [SIII] lines, are consistent with the conclusion that S/O remains constant as O/H varies over a large range in metallicity. The scatter, due



**Fig. 2.** The relationship between S/H and O/H abundances. Filled circles correspond to S/H obtained using the direct method from Te[SIII]; open circles are S/H derived from the empirical method using the parameter  $S_{23}$ . The dashed lines are  $\pm 1\sigma$  of the solar abundance ratio (Lodders 2003) shown by the solid line.



**Fig. 3.** The S/H abundance versus the [SIII]/[SII] ratio obtained for different locations within an individual galaxies; each galaxy is coded by a different symbol;  $1\sigma$  is a representative error.

mainly to observational errors, in S/O (at a given O/H) is  $\pm 0.3$  dex, still large to constrain the degree of variation in S/O expected from popIII nucleosynthesis models.

*Acknowledgements.* C.K wishes to acknowledge CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico - Brazil) for a studentship and the CSIC for a I3P postgraduate fellowship. This research was partially funded by project AYA2004-08260-C03-02 of the Spanish PNAYA.

## References

- Garnett, D.R. 1992, AJ, 103, 1330
- Garnett, D.R. 1989, ApJ, 345, 282
- Kehrig, C. et al. 2005, A&A, in preparation
- Kehrig, C. et al. 2004, AJ, 128, 1141
- Kehrig, C. 2003, Master Dissertation
- Lodders, K. 2003, ApJ, 591, 1220L
- Nomoto, K. et al. 2004, Mem. S.A.It., 75, 312
- Pérez-Montero, E. et al. 2005, in preparation
- Vilchez, J.M. & Esteban, C. 1996, MNRAS, 280, 720