



Magnetic reconnection driven by a synthetic photospheric velocity field

B. Viticchié, D. Del Moro, F. Berrilli, and A. Egidi

Dipartimento di Fisica, Università di Roma "Tor Vergata" Via della Ricerca Scientifica 1,
00133 Roma, Italy
e-mail: viticchie@roma2.infn.it

Abstract. We present a dynamic model of atmospheric magnetic field in which magnetic loop footpoints are advected by a velocity field computed using a simple n -body simulation and reproducing large spatial organization scales (e.g. mesogranulation). In this model, the advection of passive magnetic elements triggers multiple reconnection processes forcing magnetic field reconfigurations and ensuing fluctuations of total magnetic energy. The simulations reproduce a system with scale-free properties, provides frequency distribution functions for emitted magnetic energies, described by a power-law index $\alpha \sim 2.4$, and also frequency distribution for quiescent times between different energy emissions.

1. Introduction

Solar flares are energetic events in the solar atmosphere. A single flare can release a great amount of energy ($\sim 10^{30}$ erg). The released energy is stored in the atmospheric magnetic field, which, via magnetic reconnections and rapid reconfigurations, is able to emit magnetic energy explosively in very short time intervals (few min).

The frequency distribution of the peak flare energies obeys a power-law over several orders of magnitude with a robust power-law index. The distribution of quiescent times between different flares obeys time-dependent Poisson statistics for short intervals while for long intervals a power-law tail is found; this kind of distribution is well reproduced by a Levy function (Wheatland, 2002; Lepreti et al., 2001).

Here we propose a dynamical model for the atmospheric magnetic field which is repre-

sented by circular loops. Each loop is anchored to a two-dimensional domain representing the photosphere. Two footpoints define where each magnetic loop enters the photosphere. Footpoints are transported by a velocity field which presents correlations in space and time. In fact, velocity fields produce meso-granular flows by the interaction of small-scale, short-lived flows that represents granular flows. Loops are driven by the movement of footpoints, moreover they can interact via magnetic reconnection. Multiple magnetic reconnections cause magnetic field configurations to change, from this follows a variation of the magnetic energy of the system.

2. Our model

We briefly describe the dynamic model for the photospheric velocity field defined on the basis of the model from Rast (2003).

The model realizes the advection-interaction

Send offprint requests to: B. Viticchié




Fig. 1. A snapshot of the simulation from two different viewing angles. The color of the loops is proportional to th




Fig. 2. Frequency distribution for released magnetic energies. Error bars have the dimension of the dots.

between downflows in a $L \times L$ domain ($L = 256$ pixel) with periodic boundary conditions. We start our simulation with a fixed number, $N = 1000$, of downflows randomly located in the domain. The interaction (attraction) velocity between two downflows follows an exponential law $v = v_0 \cdot \exp \frac{-r}{\sigma}$ (Rast , 1995, 2003). The decay length for the interaction, σ , is a free parameter. The velocity of a downflow is defined as the sum over all the velocity due to the interactions with other downflows. Two downflows that are a unit of length (1 pixel) from each other merge to create a single down-

flow. In this process we define a new downflow placed in the same position of the strongest downflow. The intensity of the new downflow is defined as the sum of the intensity of the merging downflows. We suppose that where the downflow is located we have a vertical (orthogonal to the domain) flow velocity equal to v_0 . We also suppose that the sum over all the vertical velocities is always equal to the initial sum which is equal to 1000 flow units. From a statistical analysis (Berrilli et al. , 2003) of downflow baricenter clustering results a granulation length scale $m_G \sim 20$ pixel and




Fig. 3. Frequency distribution for quiescent time between magnetic energy emissions. Error bars have the dimension of the dots. Solid line represents the exponential fit; dashed line represents the linear fit.

a mesogranular length scale $m_M \sim 82$ pixel. The ratio of the length scales is $\frac{m_M}{m_G} \sim 5$ in agreement with the observations.

The dynamic model for the magnetic reconnection was defined referring to a model from Hughes et al. (2003). The magnetic field is represented by semicircular loops anchored by two footpoints, of opposite polarity, on the same domain $L \times L$ used for the photospheric velocity field. The photospheric horizontal velocity field moves the footpoints driving the magnetic field loops that can interact and cause magnetic reconnections, sometimes multiple ones. In agreement to the model from Hughes et al. (2003) we assume the magnetic energy of a single loop proportional to its length. Therefore, the variation of the total magnetic energy of the system, due to a reconnection process, is proportional to the variation of the sum over all loops lengths. A snapshot of a simulation run is reported in Fig. 1. In the left panel the synthetic velocity field is shown with superimposed the projection of the magnetic loops; the right panel shows a 3-d view that highlights the instantaneous magnetic con-

figuration.

With respect to the original model (Hughes et al., 2003) we implemented new features: *i*) we used the photospheric velocity field (ref. section 2) presenting correlation in space and time in agreement with the observations; *ii*) boundary conditions are periodic; *iii*) in our model loops interact at finite distance. Simulations start in a random configuration with a fixed number of magnetic loops and admit two types of magnetic interactions.

1. interaction between footpoints on the $L \times L$ domain. Footpoints with opposite polarity, from two different loops, can annihilate when they are a unit of length (1 pixel) from each other. After the interaction a new loop is defined between the two remnant footpoints of opposite polarity. Footpoints with the same polarity can aggregate in the same position in the domain.
2. interaction between loops over the $L \times L$ domain. The interaction starts when the distance between the loops is less than or equal to 1 pixel. This interaction causes a reconfiguration of the magnetic field between the involved footpoints.

Magnetic interactions, instantaneous in our model, always cause a release of magnetic energy. If more interactions can take place simultaneously, we start the reconnection process from the interaction that causes the greater emission of magnetic energy.

In our simulations, reproducing the emergence of small scale magnetic fields with a constrain on the minimum dimension for magnetic loops, loops are introduced with a fixed rate during the evolution.

The aim of the simulations is to investigate computed frequencies distributions both for energy emission and for quiescent times between emissions.

3. Statistical analysis and conclusions

The statistical analysis of magnetic energy emissions over 8 simulations brings us the results presented in Fig. 2. More in detail, the frequency distribution is well approximated by a power-law over three decades of energy, the index is equal to 2.4 ± 0.1 in agreement with observations of nanoflares (Krucker and Benz, 1998; Parnell and Jupp, 1999).

In Fig. 3 we report the frequency distribution for quiescent times. For small time interval the distribution is well approximated by an exponential function. For times greater than 10 min we found a deviation from the exponential behavior and the distribution seems well-matched

by a power-law. The fit, done on the last three points, needs to be extended in time simulation to define a statistical behavior for large quiescent times.

Our approach indicates that this class of dynamical models including space-time correlations in the photospheric velocity field can reproduce both energy and waiting time frequency distributions of nanoflares. Fairly sophisticated numerical simulations may be necessary in order to improve our model and confirm our insights.

References

- Berrilli, F., Del Moro, D., Russo, S., Consolini, G., & Straus, Th. 2005, *Astrophys. J.* Preprint doi 10.1086/432708'
- Hughes D., Paczusi M., Dendy R. O., Helander P., & Mc Clements K. G. 2003, *Phys. Rev. Letters*, 90, 131
- Krucker, S., & Benz, A. O. 1998, *Astrophys. J.*, 501, 213
- Lepreti, F., Carbone, V., & Veltri, P. 2001, *Astrophys. J.*, 555, 133
- Parnell, C. E., & Jupp P. E. 1999, *Astrophys. J.*, 529, 554
- Rast, M. P. 1995, *Astrophys. J.*, 443, 863
- Rast, M. P. 2003, *Astrophys. J.*, 597, 1200
- Van Siclen, C. 1997, *Phys. Rev.*, E 56, 5211
- Wheatland, M. S., & Litvinenko, Y. E. 2002, *Solar Phys.*, 211, 255