



The chemical composition of Cepheids in the Milky Way and the Magellanic Clouds

M. Mottini¹, M. Romaniello¹, F. Primas¹, G. Bono²,
M.A.T. Groenewegen³, and P. François⁴

¹ European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching bei München, Germany

² INAF-Osservatorio Astronomico di Roma, via Frascati 33, I-00040 Monte Porzio Catone, Italy

³ Instituut voor Sterrenkunde, Celestijnenlaan 200B, B-3001 Leuven, Belgium

⁴ Observatoire de Paris-Meudon, GEPI, 61 avenue de l'Observatoire, F-75014 Paris, France

Abstract. We have measured the elemental abundances of 68 Galactic and Magellanic Cepheids from FEROS and UVES high-resolution and high signal-to-noise spectra in order to establish the influence of the chemical composition on the properties of these stars (see Romaniello et al. 2005). Here we describe the robust analytical procedure we have developed to accurately determine them. The resulting iron abundances span a range between ~ -0.80 dex for stars in the Small Magellanic Cloud and $\sim +0.20$ dex for the most metal-rich ones in the Galaxy. While the average values for each galaxy are in good agreement with non-pulsating stars of similar age, Cepheids display a significant spread. Thus it is fundamental to measure the metallicity of individual stars.

Key words. Stars: abundances – Cepheids

1. Introduction

Cepheid stars, through their Period-Luminosity (PL) relation, are one of the key tools in determining the extragalactic distance scale. In spite of its fundamental importance, to this day, it is still unclear the role played by the chemical composition on the pulsational properties of these stars. Recent theoretical works give different results: some authors claim that metallicity has negligible effects on the PL relation (Chiosi et al. 1992; Sandage et al. 1999; Baraffe & Alibert 2001); others claim that there is a significant depen-

dence of the PL relation on metallicity (Bono et al. 1999; Fiorentino et al. 2002; Marconi et al. 2005).

Although most of the observational efforts aiming at deriving the chemical composition of Cepheids and its effects on PL relation have used indirect means, like the measurement of oxygen abundances in the H II regions (e.g. Sasselov et al. 1997; Kennicutt et al. 1998; Sakai et al. 2004), some of them have focused on a direct chemical analysis (Fry & Carney 1997; Luck et al. 1998; Andrievsky et al. 2002; Luck et al. 2003). Our study follows this second approach: we analysed a sample of high resolution spectra and derived α -elements and

Send offprint requests to: M. Mottini

iron abundances spectroscopically. The preliminary results for a sub-sample of 37 stars have already been presented in Romaniello et al. (2005). Here we present the complete sample.

2. The data sample

We used the following selection criteria to choose the Cepheids of this data set: stars from three galaxies (Milky Way, LMC and SMC) in order to cover a factor of about ten in metallicity, well-determined intensity-mean magnitude in B, V and K bands for all the Cepheids (Laney & Stobie 1994; Groenewegen et al. 2004; Storm et al. 2004); a wide range of periods in order to cover the entire PL relation (from 2 to 98 days).

We have a sample of 68 Cepheids: 32 Galactic Cepheids, 22 LMC and 14 SMC Cepheids. The spectra of the Galactic Cepheids have been collected with FEROS (R=48 000, spectral coverage: 3700-8600 Å) at the ESO 1.5 meter telescope (La Silla) with a $S/N \sim 150$ for $\lambda > 5000$ Å, $S/N \sim 70$ for $\lambda > 4000$ Å. The spectra of the Magellanic Cepheids have been collected with UVES (R=40 000, spectral coverage: 4800-6800 Å for the red arm) at the ESO VLT-Kueyen (Paranal) with a $S/N \sim 70$ for $\lambda > 5800$ Å, $S/N \sim 50$ for $\lambda > 4800$ Å.

3. The analysis

In order to derive elemental abundances one needs to know the stellar parameters of the star and have a reliable line list available.

As a first step we carefully assembled the line list. We have assembled our Fe I-Fe II and α -elements line list from Clementini et al. (1995), Fry & Carney (1997), Kiss & Vinko (2000) and Andrievsky et al. (2002) including a selection of lines from VALD (Kupka et al. 1999; Ryabchikova et al. 1999). The VALD lines have been selected for effective temperatures typical of Cepheid stars (4500-6500 K). We have, then, visually inspected each line on the observed spectra, in order to check their profile and to discard blended lines. In order to do so, we have searched the VALD database (with the command *extract stellar*) for all the

listed lines between 4800 and 7900 Å on stellar spectra characterised by parameters typical of Cepheids ($T_{\text{eff}}=4500$ K, 5500 K and 6500 K, $\log g=2$ and $v_t=3$ kms⁻¹). We then overplotted all the lines found in VALD, that fall within ± 3 Å from each of our lines, and checked for their possible contribution to the equivalent width of our line. Any contribution larger than 5% of the line strength made us discarding the line under scrutiny.

Our final list includes 246 Fe I lines and 17 Fe II lines spanning a spectral range from 4800 Å to 7900 Å which is the range covered by our FEROS spectra. For the UVES spectra this list is slightly reduced due to their narrower spectral range. Simultaneously we also looked at α -elements (O, Na, Mg, Al, Si, Ca and Ti) and we have assembled a list of almost 200 lines. For all the lines we have adopted the physical properties (oscillator strengths, excitation potentials) listed in VALD.

As a second step we measured the equivalent widths (EW) of all the lines assembled as described above. Because of their large number, we have used a semi-interactive routine developed by one of us (PF, *fitline*). This code is based on genetic algorithms, which mimic how DNA can be affected to make the evolution of species (Charbonneau 1995) and it uses a Gaussian fit. For about 15% of the lines, the Gaussian profile adopted by *fitline* could not satisfactorily reproduce the observed profile. For these cases (usually very broad or asymmetric lines), the equivalent widths were measured manually with the IRAF *splot* task. The mean difference, as computed for those lines for which both methods could be applied, is around 1.5 mÅ, comparable to the average error on the EW inferred from the quality of the data (Equation 7, Cayrel 1988). The outcome of this test is that we can then safely use all our EW measurements, independently of the method followed to derive them.

With these EWs in hand we then derived the microturbulent velocity and gravity of the studied star. We used as first guess values typical of Cepheids as inferred from previous studies (see Section 1 for references) then we got the final constraints from the abundance analysis: from the minimization of the $\log(\text{Fe}/\text{H})$ vs




Fig. 1. Histograms of the $[\text{Fe}/\text{H}]$ ratios derived for all the stars of our sample in the Galaxy, the LMC and the SMC. The solid lines indicate the mean values found using F and K supergiants by Hill et al. (1995) and Hill (1997).

EW slope for the microturbulent velocity and from the ionization balance for the $\log g$. For the determination of the effective temperature we used the line depth ratios method described in Kovtyukh & Gorlova (2000).

Finally we determined the elemental abundances of our stars by using the Kurucz WIDTH9 code (Kurucz 1993) and LTE model atmospheres computed without the overshooting approximation and with the most recent opacity distribution functions (Castelli & Kurucz 2003).

4. Results and future developments

Figure 1 shows the distributions of the iron abundances of our programme stars in the Galaxy and the Magellanic Clouds. In the latter case we compare our results with the mean values obtained by Hill et al. (1995) and Hill (1997) for F and K supergiants in the LMC and SMC, respectively. We note a good agreement with our mean values of the iron content. For our Galactic sample, we find that the mean value of the iron content is solar ($\sigma = 0.10$), with a range of values between -0.18 dex and $+0.25$ dex. These two extremes are represented respectively by 2 and 1 stars (out of the 32

in total). We note that more than 50% of the stars have a sub-solar metallicity. For the LMC sample, we find a mean metallicity value of ~ -0.33 dex ($\sigma = 0.13$), with a range of values between -0.62 dex and -0.10 dex. Here, the more metal-rich extreme is just an isolated case, while the metal-poor end of the distribution is represented by 3 stars. For the SMC sample, we find that the mean value is about ~ -0.75 dex ($\sigma = 0.08$), with a range of values between -0.87 and -0.63 . While the average values for each galaxy are in good agreement with non-pulsating stars of similar age, Cepheids display a significant spread. Thus it is fundamental to measure the metallicity of individual stars. Detailed discussion of these results and their comparison with previous studies will be presented in a forthcoming paper (Mottini et al. 2006), along with an analysis of the effects of the metallicity on the Cepheid Period-Luminosity relation, also showed by Romaniello et al. in these proceedings. In the very near future we will focus on the analysis of the α -elements results and their interpretation.

References

- Andrievsky, S.M. et al. 2002, *A&A*, 381, 32
 Baraffe, I. & Alibert, Y. 2001, *A&A*, 371, 592
 Bono, G. et al. 1999, *ApJ*, 512, 711
 Castelli, F. & Kurucz, R.L. 2003, in *Modelling of Stellar Atmospheres*, 210th IAU Symposium, Edited by N. Piskunov, W.W. Weiss and D.F. Gray. Published by the Astronomical Society of the Pacific, p.A20
 Cayrel, R. 1988, in *The Impact of Very High S/N Spectroscopy on Stellar Physics*, 132th IAU Symposium, Kluwer Academic Publishers, Dordrecht., p.345
 Charbonneau, P. 1995, *ApJSS*, 101, 309
 Chiosi, C. et al. 1992, *ApJ*, 387, 320
 Clementini, G. et al. 1995, *AJ*, 110, 2319
 Fiorentino, G. et al. 2002, *ApJ*, 576, 402
 Fry, A.M. & Carney, B.W. 1997, *AJ*, 113, 1073
 Groenewegen, M.A.T. et al. 2004, *A&A*, 420, 655
 Hill, V. 1997, *A&A*, 324, 435
 Hill, V., Andrievsky, S.M. & Spite, M. 1995, *A&A*, 293, 347
 Kennicutt, R.C. et al. 1998, *ApJ*, 498, 181
 Kiss, L.L. & Vinko, J. 2000, *MNRAS*, 314, 420
 Kovtyukh, V.V. & Gorlova, N.I. 2000, *A&A*, 351, 597
 Kupka, F. et al. 1999, *A&AS*, 138, 119
 Kurucz, R.L. 1993, *CD-ROMS #1*, 13, 18
 Laney, C.D. & Stobie, R.S. 1994, *MNRAS*, 226, 441
 Luck, R.E. et al. 1998, *AJ*, 115, 605
 Luck, R.E. et al. 2003, *A&A*, 401, 939
 Marconi, M., Musella, I. & Fiorentino, G. 2005, *ApJ*, 632, 590
 Mottini, M. et al. 2006, to be submitted to *A&A*
 Romaniello, M. et al. 2005, *A&A*, 429, L37
 Ryabchikova T.A. et al. 1999, *proc. of the 6th International Colloquium on Atomic Spectra and Oscillator Strengths*, Victoria BC, Canada, 1998, *Physica Scripta T83*, 162
 Sakai, S. et al. 2004, *ApJ*, 608, 42
 Sandage, A., Bell, R.A. & Tripicco, M.J. 1999, *ApJ*, 522, 250
 Sasselov, D.D. et al. 1997, *A&A*, 324, 471
 Storm, J. et al. 2004, *A&A*, 415, 531