



# First unambiguous asteroseismologic modelling of a $\gamma$ Doradus star

A. Moya<sup>1</sup>, A. Grigahcene<sup>2</sup>, J.C. Suárez<sup>3</sup>, S. Martín-Ruiz<sup>3</sup>, P.J. Amado<sup>4</sup>,  
M.A. Dupret<sup>1</sup>, R. Garrido<sup>3</sup>, and E. Rodríguez<sup>3</sup>

<sup>1</sup> Observatoire de Paris, LESIA, UMR 8109, 92195 Meudon, France

<sup>2</sup> CRAAG, Algiers Observatory BP 63 Bouzareah 16340, Algiers, Algeria

<sup>3</sup> Instituto de Astrofísica de Andalucía-CSIC, Granada CP. 3004, Spain

<sup>4</sup> European Southern Observatory, Alonso de Cordova 3107, Santiago 19, Chile

**Abstract.** An asteroseismologic model of the  $\gamma$  Dor star 9 Aurigae is discussed in this work. This is the first time a complete asteroseismologic scheme for analyzing  $\gamma$  Dor stars is applied. From standard photometric observations (at least three oscillation frequencies, a photometric error box in the HR diagram and multicolor photometric observations, in this case in the Strömngren system) we can reduce the possible theoretical models for this star, providing constraints to the stellar parameters as mass, overshooting, metallicity, MLT parameter  $\alpha$ , Brunt-Väiälä integral ( $I_{th}$ ), etc. Simultaneously, an estimate of the modal identification of the observed frequencies is also obtained.

This can be possible by the application, for the first time, of a complete procedure where different theoretical and computational techniques, recently developed, are linked and compared with photometric observations. The Frequency Ratio Method (FRM) and the Time Dependent Convection (TDC) theory are the basis of this complete procedure.

**Key words.** Stars:  $\gamma$  Dor – Stars: 9 Aurigae – Stars: Modeling

## 1. Introduction

The principal achievement of asteroseismology is to improve our knowledge of the stellar interiors by using additional observational constraints: the oscillation frequencies. To do so, the first step to be done is to identify the modes these frequencies belong to. This will make it possible to better define the theoretical model fitting all the observational constraints, giving information about the accuracy of the theories describing the star and the ad hoc parameters some theories have.

With the recent development of some techniques for comparing theoretical predictions with observations, mainly the Frequency Ratio Method (FRM) (Moya et al. 2005; Suarez et al. 2005) and the Time Dependent Convection (TDC) (Grigahcène et al. 2005; Dupret et al. 2005; Dupret et al. 2005), we are now able to construct a self-consistent procedure to test theoretical predictions maximizing the constraints. This procedure, applied to the  $\gamma$  Dor star 9 Aurigae gives as a result a single possible model fitting all observations within the present theoretical descriptions.

---

Send offprint requests to: A. Moya (moya@iaa.es)

**Table 1.** 9 Aurigae model characteristics

Mass	$T_{eff}$	$\log g$	$\log \frac{L}{L_{\odot}}$	$\log \frac{R}{R_{\odot}}$	Age (My)	[Fe/H]
1.4	7006	4.28	0.63	1.41	600	-0.1

**Table 2.** 9 Aurigae theoretical characteristic and modal identification

$\alpha_{MLT}$	$I_{th}$	$\alpha_{ov}$	$n_1$	$n_2$	$n_3$	$\ell$
1.6	681.5	0.3	57	59	133	2

## 2. Self-consistent procedure

The minimum observational requirements needed to apply this procedure are 1) a photometric error box in the HR diagram, and 2) at least three pulsational frequencies observed in multicolor photometry. Nevertheless spectroscopic information will be also very useful. The starting point is the photometric error box and the three frequencies. For 9 Aurigae the observational characteristics and observational frequencies ( $f_1 = 0.7948$  cycle  $d^{-1}$ ,  $f_2 = 0.7679$  cycle  $d^{-1}$  and  $f_3 = 0.3429$  cycle  $d^{-1}$ ) are obtained from Zerbi et al. (1997).

The FRM gives a first restriction in modal identification, overshooting, metallicity, mass and  $T_{eff}$ . Taking into account only models given by the FRM we can study the instability of the theoretical frequencies with TDC, providing in addition, predictions for the multicolor photometric observables. Comparing these predictions with the observations the number of possible models fulfilling all the observational constraints is highly reduced. In this case only one model from our grid has been selected (see Tabs 1 and 2).

## 3. Problems

Some problems has been encountered: 1) The Strömgren filters ratio  $u/y$  cannot be reproduced by the models, 2) the frequency  $f_3$  is

never predicted overstable and 3)  $\alpha_{ov}$  is larger than expected.

Nevertheless the obtaining of only one theoretical model fulfilling all the observational constraints make it possible to test the accuracy of the theory used.

*Acknowledgements.* The authors acknowledges the financial support of the the European Marie Curie action MERG-CT-2004-513610 and the Spanish Plan Nacional del Espacio under project ESP2004-03855-C03-01.

## References

- Dupret, M.-A., Grigahcène, A., Garrido, R., Gabriel, M., & Scuflaire, R. 2005, A&A, 435, 927
- Dupret, M.A., Grigahcène, A., Garrido, R., De Ridder, J., Moya, A., Suárez, J. C., Scuflaire, R., Gabriel, M. Goupil, M.J., 2005, this Meeting.
- Grigahcène, A., Dupret, M.-A., Gabriel, M., Garrido, R., & Scuflaire, R. 2005, A&A, 434, 1055
- Moya, A., Suárez, J. C., Amado, P. J., Martin-Ruíz, S., & Garrido, R. 2005, A&A, 432, 189
- Suárez, J. C., Moya, A., Martin-Ruíz, S., Amado, P. J., Grigahcène, A., & Garrido, R. 2005, Accepted in A&A.
- Zerbi, F. M., et al. 1997, MNRAS, 290, 401