

Hybrid stars with a simultaneous δ Scuti and γ Doradus stars pulsational behaviour

A. Grigahcène^{1,2}, S. Martín-Ruiz², M. A. Dupret³, R. Garrido² and M. Gabriel⁴

¹ Algiers Observatory, CRAAG, Algiers, Algeria

² Instituto de Astrofísica de Andalucía-CSIC, Granada, Spain

³ Observatoire de Paris-Meudon, LESIA, France

⁴ Institut d'Astrophysique et de Géophysique de l'Université de Liège, Belgium e-mail: ahmed@iaa.es

Abstract. Computations of theoretical instability strips for δ Scuti (Sct) and γ Doradus (Dor) stars are presented. The time-dependent convection (TDC) theory of Gabriel (1996) and Grigahcène et al. (2005) is adopted in our models. The theoretical Instability Strips (IS) obtained for δ Sct and γ Dor stars overlap indicating a possibility of hybrid stars showing simultaneously both γ Dor IS and δ Sct excited modes. Models with unstable long periods gravity modes (γ Dor type) show also unstable short periods pressure modes (δ Sct type). First example of such a star is given by Henry & Fekel (2005).

Key words. Stars: oscillations – Convection – Stars: interior – Stars: variables: general

1. Introduction

Because of their location in the HR diagram, early when the γ Dor new class of pulsating stars was discovered, the question arised of the relationship between them and the δ Sct stars (Breger & Beichbuchner 1996). Search of stars pulsating simultaneously with δ Sct p-modes and γ Dor g-modes have been done by Handler & Shobbrook (2002) and recently by Henry & Fekel (2005).

2. Theoretical models

Fig. 2 shows theoretical IS for γ Dor (dashed lines) for $\ell = 1$ modes and δ Sct red edge (dashed-pointed line) for the fundamental radial mode, computed with our TDC non-

adiabatic models with $\alpha = 1.8$. Asterix symbols and filled squares are observed positions of bona fidae γ Dor stars given by Handler (2002) and Martín (2003a) respectively. Dots represent the observed positions of δ Sct stars from the catalogue of Rodríguez, López-Gonzalez & López de Coca (2000). A significant part of the γ Dor IS is inside the δ Sct one. All the theoretical models in this intersection have both types of unstable modes. Existence of hybrid δ Sct - γ Dor is predicted by our theoretical models (Grigahcène et al. 2005). Fig. 1 gives the damping rate (negative for excited modes) as a function of period (en hours). It shows clearly that models with unstable long periods g-modes have also unstable short periods p-modes. These short period unstable modes are the typical δ Sct type modes.

Send offprint requests to: A. Grigahcène




Fig. 1. Simultaneous excitation of g and p-modes for the same model: $M = 1.6M_{\odot}$, $\alpha = 1.8$ and $T_{eff} = 6995$ K.

3. Systematic observational search

First example of a star pulsating with both γ Dor and δ Sct frequencies is given by Henry & Fekel (2005) (circled plus in Fig. 2). It shows periods: $0.0484 \text{ d} < \text{period} < 0.4033 \text{ d}$. It lays in the intersection between theoretical γ Dor and δ Sct IS. A systematic search for the γ Dor in 10 open clusters was performed Martín (2003a) by using uvby photometry. From these observations, eight new γ Dor were found (filled squares Fig. 2). On the light of the TDC theoretical results, a new Fourier analysis has been made with the goal of detecting high frequencies in their frequency spectra. Two codes have been used: Multifre (Martín 2003b) and the same method as Rodríguez et al. (1998). No δ Sct type frequencies have been found in these stars.

4. Conclusions

We conclude that there are no theoretical arguments against the existence of hybrid δ Sct - γ Dor variables. It is very important to continue the search of such stars (even a first discovery has been done), which would be of considerable interest for asteroseismology: high order g-modes allow probing the deep interior while p-modes allow probing the more superficial layers.




Fig. 2. Theoretical IS for γ Dor (dashed lines) and δ Sct red edge (dashed-pointed line), compared with observations of *bona fide* γ Dor stars (*) of Handler (2002) and (■) of Martín (2003a) and observations of δ Sct stars (○) of Rodríguez, López-Gonzalez & López de Coca (2000). (⊕) is the hybrid star of Henry & Fekel (2005).

References

- Breger, M. & Beichbuchner, F., 1996, A&A, 313, 851
 Gabriel, M., 1996, Bull. Astron. Soc. of India, 24, 223
 Grigahcène, A., Dupret M.-A., Gabriel M., Garrido R. et al., 2005, A&A, 434, 1055
 Handler, G. & Shobbrook, R. R., 2002, MNRAS, 333, 251
 Handler, G., 2002, <http://www.astro.univie.ac.at/~dsn/gerald/gdorlist.html>
 Handler, G., Balona, L. A., Shobbrook, R. R., et al., 2002, MNRAS, 333, 262
 Henry, G. W. & Fekel, F. C., 2005, Astron. J., 129, 2026
 S. Martín, 2003a, in ASP Conf. Ser. 292: Interplay of Periodic, Cyclic and Stochastic Variability in Selected Areas of the H-R Diagram, 59
 Martín et al., 2003b, A&A, 401, 1077
 Rodríguez et al., 1998, A&A, 338, 905
 Rodríguez, E., López-Gonzalez, M. & López de Coca, P., 2000, A&AS 144, 469