



# The O I $\lambda 8446$ emission line in NLS1 galaxies

V. Botte, S. Ciroi, F. Di Mille, and P. Rafanelli

Department of Astronomy, University of Padova, vicolo dell'Osservatorio 2, I- 35122 Padova, Italy

**Abstract.** In studying Ca II  $\lambda\lambda 8498, 8542, 8662$  spectral region of NLS1, we have observed that 7 NLS1 do not show O I  $\lambda 8446$  line emission. We have investigated the reasons of this disappearing analysing the excitation mechanism of O I  $\lambda 8446$  and the physical conditions of BLR that permit its emission. With a preliminary study of BLR properties we have found that O I  $\lambda 8446$  emission required a density  $n_H$  between  $10^{11}$  and  $10^{12}$  and ionization parameter  $U$  between  $-1.8 \div -3$  in logarithm.

## 1. Introduction

The origin and configuration of the gas which emits broad lines in type 1 AGN is not clear yet, especially in NLS1. O I  $\lambda 8446$  usually the strongest O I line in the optical region, has been broadly studied in a large number of objects. In type 1 AGN, this emission line is very common. In particular its excitation mechanisms of O I  $\lambda 8446$  have been investigated by many different authors especially because investigating the O I line formation mechanisms would also provide the knowledge about physical properties of the FeII emitting region. We have focused our attention on a feature showed by NLS1.

## 2. The disappearing of O I $\lambda 8446$ line in some NLS1

We have collected from literature (Persson S. E. 1988; Rodriguez-Ardila et al. 2002; Botte et al. 2005) a sample of thirty nuclear spectra of NLS1 covering the range in which the

O I  $\lambda 8446$  falls, and we have found for seven of them the absence of this line.

Our aim is to try to establish why O I  $\lambda 8446$  emission is not present in all NLS1 galaxies in order to explore the physics of the gas emitting their spectrum and compare it to that one of the Seyfert 1 (S1).

Firstly we have carried out photoionization model calculations by using the code CLOUDY version 94.00 (Ferland 1996, 2000) for the BLR. We have made several models covering the ranges  $-3 \div -1$  for the ionization parameter ( $U$ ) and  $8 \div 12$  for the hydrogen density of the gaseous clouds ( $n_H$ ) in logarithm scale; solar abundances were adopted according to Grevesse and Anders (1989), and for the shape of ionizing continuum a power law with spectral index ( $\alpha$ ) in the range  $-2 \leq \alpha \leq -1$  was assumed.

As final step we have plotted the theoretical values of  $\lambda 8446/H\beta_{broad}$  ratios as a function of  $U$ ,  $n_H$  and  $\alpha$ . On these plots we have marked the region covered by the measured values of  $\lambda 8446/H\beta_{broad}$  obtained from a sample of 10 NLS1. Excluding the upper and the lower values the  $\lambda 8446/H\beta_{broad}$  ratios fall in the range  $0.17 \div 0.30$  implying  $n_H$  between

---

Send offprint requests to: V. Botte

**Table 1.** Narrow-Line Seyfert 1

Name	R.A.(J2000) hh mm ss	Dec.(J2000) ° ' "	z
Mrk359	01 27 32.5	+19 10 44	0.0167
IR04321+4008	04 34 41.5	+40 14 22	0.0201
CTS H34.06	06 09 17.5	-56 06 58	0.0318
CTS J03.19	10 15 56.6	-20 02 27	0.0547
CTS J04.08	10 31 57.3	-18 46 34	0.0403
IC3599	12 37 41.2	+26 42 27	0.0221
Kaz163	17 46 59.8	+68 36 39	0.0637

$10^{11}$  and  $10^{12}$  and  $\log U$  between  $-1.8 \div -3$ . In order to test our model, we have constructed the same graphic for other two broad emission line, He II 4686 and H $\alpha$ . Combining all results, we have obtained  $n_H \sim 10^{11} \text{ cm}^{-3}$ ,  $\log U$  between  $-1.8 \div -2.6$  and  $\alpha$  between  $-1.75 \div -1.5$ .

### 3. Results and conclusions

NLS1 are a subgroup of S1 galaxies that display very peculiar properties which are not yet completely understood.


We have added to these another peculiarity: some NLS1 galaxies do not show O I  $\lambda 8446$  emission in their spectrum.

With a preliminary study of BLR properties we have found that O I  $\lambda 8446$  emission requires density  $n_H$  between  $10^{11}$  and  $10^{12} \text{ cm}^{-3}$  and ionization parameter U between  $-1.8 \div -3$  in logarithm. This result is in agreement with Kuraszkiwicz et al. (2000) conclusion: in NLS1 objects the BLR clouds have lower ionizing parameters and higher densities than normal AGNs.

As a consequence, a possible explanation for the absence of O I  $\lambda 8446$  in some objects, could be that the physical conditions of BLR do not allow the production of this line by Ly $\beta$  fluorescence (a partially ionized region in the clouds emitting Ly $\beta$  photons, O I atoms and large optical depth of H $\alpha$ .)

Another possibility could be the presence of optically thin gas in the BLR of NLS1.

The disappearing of O I  $\lambda 8446$  lines in some NLS1 remain an open question but it is clear that in order to understand the cause of this unobserved lines our models need to be improved, performing new ones in individual



**Fig. 1.** O I  $\lambda 8446$  relative intensity versus U, for different densities; continuous line:  $10^{12} \text{ cm}^{-3}$ ; large dashed lines:  $10^{11} \text{ cm}^{-3}$ ; dashed lines:  $10^{10} \text{ cm}^{-3}$ ; small dashed lines:  $10^9 \text{ cm}^{-3}$ . The regions marked by red colour correspond to O I  $\lambda 8446$  relative intensity measured.

NLS1 with observations of a larger spectral range.

### References

- Botte V., Ciroi S., Di Mille F., Rafanelli P. & Romano A. 2005, AJ, 357, 789  
 Ferland G. J., 1996, HAZY, A Brief Introduction to CLOUDY (Lexington: Univ Kentucky Dept. Phys. Astron.)  
 Ferland G. J., 2000, Rev. Mexicana Astron. Astrofis. Ser. Conf., 9, 153  
 Grandi S. A., 1980, ApJ, 238, 10  
 Kuraszkiwicz J., Wilkes B. J., Czerny B. & Mathur S. 2000, ApJ, 542, 692  
 Mathur, S. 2000, MNRAS, 314, 17  
 Pounds, K. A., Stanger, V. J., Turner, T. J., King, A. R., & Czerny, B. 1987, ApJ, 330, 751  
 Rodriguez-Ardila A., Viegas S. M., Pastoriza M. G., Prato L., Donzelli C. L., 2002, ApJ, 572, 94