

# Three years monitoring of 3C 273 with INTEGRAL and XMM-Newton

V. Bianchin<sup>1</sup>, P. Grandi<sup>1</sup>, L. Foschini<sup>1</sup>, and G. G. C. Palumbo<sup>2</sup>

<sup>1</sup> INAF/IASF-Bologna, Via Gobetti 101, 40129 Bologna, Italy  
e-mail: bianchin@iasfbo.inaf.it

<sup>2</sup> Università di Bologna, Dipartimento di Astronomia, Via Ranzani 1, 40126 Bologna, Italy

**Abstract.** We present preliminary results of the analysis of 3C 273 observed with *INTEGRAL* and *XMM-Newton* during 2003 – 2005. The available data span from optical to hard X frequencies and show significant variability. Simulations with *SIMBOL-X* are also presented.

**Key words.** quasars: individual: 3C 273 – X-rays: galaxies – Galaxies: active

## 1. Introduction

We selected all the *INTEGRAL* publicly available pointings with the source in the FCFOV ( $< 5^\circ$  to the boresight) within the period 2003–2005, resulting in a total exposure of about 1 Ms. We analyzed the IBIS data using the standard software OSA v. 5.1. We retrieved the public *XMM-Newton* data collected over the same period, for a total exposure of 128 ks and analyzed them by means of the standard software SAS v. 7.0.


Lightcurves have been produced for 3C 273 from optical (*XMM/OM*) to hard X (*INTEGRAL/ISGRI*) bands, obtained with a binning time from 1.3 to 65 ks for OM data, 3 ks for PN and 90 ks for ISGRI (Fig.1). The lightcurves in the optical and ultraviolet bands are constant within errors, while strong flux variations show up in the X-ray bands. At the beginning of 2003 the source was increasing its flux level in both 0.2 – 1 and 3 – 10 keV bands, followed by a long decrease (2004) and

a slow enhancement in 2005. The flux in both the 20 – 40 keV and 40 – 100 keV ranges suggests an anticorrelated trend to the soft X emission and a correlation to the radio flux, that suggests the presence of a jet.

Moreover the dominance of the soft X emission during the low jet contribution should reveal the Seyfert like component as suggested by Grandi & Palumbo (2004). A spectral analysis of all *XMM* observation is in progress and preliminary fits with a simple broken power-law confirm the trend found by Foschini et al. (2006).


In order to disentangle the Seyfert and jet component a spectral analysis of simultaneous data is needed. However *INTEGRAL* needs significantly longer exposure times than *XMM-Newton*. Indeed, as shown in Fig. 2, with 4 ks of exposures, it is possible to achieve only a low significance detection in two energy bands, despite 3C 273 is one of the brightest blazars in the sky.

The *SIMBOL-X* mission is expected to guarantee a high sensitivity in the hard X-rays




**Fig. 1.** From the top to bottom panel: *XMM*/OM lightcurves for filters V, B, U, UVW1, UVM2, UVW2; *XMM*/PN lightcurves in the 0.2 – 1 and 3 – 10 keV bands; *INTEGRAL*/ISGRI lightcurves in the 20 – 40 and 40 – 100 keV bands.

band. Simulations of observation of 3C 273 show that the different spectral components can be separated already with a 1 hour expo-



**Fig. 2.** Quasi-simultaneous spectral fit of *XMM-Newton* and *INTEGRAL* observation of 3C 273 performed on 2005 – 07 – 10. See text for more details.



**Fig. 3.** Simulation of 1 hour observation of 3C 273 with *SIMBOL-X*, by adopting the same spectral parameters of the fit on the July 2005 observation.

sure (Fig. 3) and parameters can be very well constrained with a 100 ks observation.

*Acknowledgements.* This work was partially supported by the Italian Space Agency (ASI) under contract I/R/046/04.

## References

- Foschini L. et al. 2006, *A&A*, 453, 829  
 Grandi P. & Palumbo G.G.C., 2004, *Science*, 306, 998