



The most powerful explosions in the Universe: genesis and evolution of supernova and Gamma-Ray Burst Italian programs at ESO

E. Pian^{1,2,3}

¹ Istituto Nazionale di Astrofisica – Osservatorio Astronomico di Trieste, Via Tiepolo 11, I-34143 Trieste, Italy

² Scuola Normale Superiore di Pisa, Piazza dei Cavalieri 7, 56126 Pisa, Italy
e-mail: elena.pian@sns.it

³ INFN-Sezione di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy

Abstract. The Italian communities engaged in Gamma-Ray Burst (GRB) and supernova research have been using actively the ESO telescopes and have contributed to improve and refine the observing techniques and even to guide the characteristics and performances of the instruments that were developed. Members of these two communities have recently found ground for a close collaboration on the powerful supernovae that underlie some GRBs. I will review the programs that have led to some important discoveries and milestones on thermonuclear and core-collapse supernovae and on GRBs.

Key words. Radiation mechanisms: general – Gamma-ray burst: general – Stars: massive – supernovae: general – Galaxies: star formation – white dwarfs

1. Introduction

Supernovae and Gamma-Ray Bursts (GRB) exhibit similar energy outputs, comparable to the binding energy of a compact star, $\sim 10^{53}$ erg, that they release in various forms, including electromagnetic radiation from radio to gamma-rays. There is evidence that both explosion types are non-spherical, although the degree of asymmetry is very different, and its effect on the observed radiation is magnified by the presence of relativistic conditions, which are absent in thermonuclear supernovae, occasionally present and mild in core-collapse supernovae and extreme in GRBs. The evolution

of the progenitor star influences the explosion, and its observable aftermath may depend on intrinsic conditions, but also on the circumstellar medium and on the viewing angle. This generates a highly diversified range of observed properties of the explosion remnants, that hold the key for the understanding of these phenomena. Selected results obtained so far on these sources with observations at ESO telescopes by Italian scientists are here reviewed.

2. Supernova programs at ESO

Five years after Italy had joined ESO, Supernova 1987A went off in the Large Magellanic Cloud, and drove a flurry of observations and theoretical studies that con-

tributed to boost the interest in supernova research in Italy and to establish systematic observing programs. Among these, two ESO Key Programs led by the group at Padua Astronomical Observatory, and involving the 3.6m, NTT, 2.2m, and Dutch telescopes, were started in the 1990s. These contributed critically to increase the statistics of well monitored supernovae and can be considered the precursors of ESO observing projects on unpredictable targets, i.e. sources whose coordinates – because of their explosive nature – cannot be known a priori. Thereafter, supernova programs have been entertained by the Padua team at ESO regularly and have had an impact on international supernova research, including the characterization of “branch-normal Type Ia supernovae” (Patat et al. 1996), of sub-luminous Ia supernovae (Turatto et al. 1996), and the contribution to work on cosmology (Hamuy et al. 1996; Perlmutter et al. 1998) that eventually led to the discovery of the accelerated expansion of the Universe (Riess et al. 1998; Schmidt et al. 1998; Perlmutter et al. 1999).

This detailed work on individual Type Ia supernovae¹ has underlined the remarkable diversity in these sources, that were traditionally considered rather homogeneous – at least photometrically – and as such used as standard candles. Supernovae Ia have been known for about 20 years not to be uniform after all, because their maximum luminosities span a magnitude or so. The relationship reported by Phillips (1993) between the maximum luminosity and the rate of light decay in the B band after maximum restores the reliability of Type Ia supernovae as standard candles, but, again, not all of them comply with this relationship. A third parameter seems to be necessary to describe the full range of Ia supernovae properties, and this has been identified as the velocity gradient, i.e. the rate at which the photospheric velocity, estimated from the blue-shift of the SiII 6355Å line, decreases with time.

In a seminal paper by S. Benetti and collaborators (2005), three sub-classes of objects

¹ a.k.a. thermonuclear supernovae, to be distinguished from core-collapse supernovae

are distinguished in a sample of 26 well monitored Ia supernovae by their location in the space of photospheric velocity decline rate and maximum luminosity. This segregation is suggested to be related to the type of explosion, i.e. deflagration versus detonation, or a transition between the two (delayed detonation). However, subsequent work on this same sample of Ia supernovae, based on both early and late photometry and spectra, showed that almost all explosions are produced by the burning of a white dwarf star of rather constant mass, $1.05 \pm 0.09 M_{\odot}$, the ones that synthesize more nuclear statistical equilibrium elements (^{56}Ni , ^{58}Ni , ^{54}Fe) producing less intermediate mass elements (e.g. silicon), and viceversa, so that the sum of their masses is constant with relatively small dispersion. This result is summarized in the flipped-Z-shaped diagram² of mass versus $\Delta m_{15}(B)$ discussed in Mazzali et al. (2007a) and it suggests that the explosion mechanism is the same in all cases (most probably delayed detonation), with the number of ignition spots determining the final nucleosynthesis.

Later on, it was suggested that the differences within the sub-classes of Ia supernovae based on photospheric velocity decline rates can be determined by a different viewing angle, implying that the explosion is asymmetric (Maeda et al. 2010), a result confirmed also by ESO spectropolarimetry (Maund et al. 2010).

3. Gamma-Ray Bursts programs at ESO

Until early 1997, GRBs were one of the biggest outstanding enigmas of astrophysics: the lack of detected counterparts at lower energies than the gamma-rays was preventing any firm conclusion about their nature and even about their distance scale, conjectures ranging from galactic to cosmological scenarios (Lamb 1997; Paczyński 1997). GRB970228 was localized with arc-minute precision by the Italian-Dutch *BeppoSAX* satellite, which made possible the detection of its X-ray afterglow, the first GRB counterpart ever detected (Costa et al. 1997).

² Sometimes called “Zorro” diagram

The afterglow era had started. The accurate localization by the *BeppoSAX* instruments of the GRB and its afterglow allowed also the unambiguous identification of the counterpart at optical wavelengths (Van Paradijs et al. 1997). The NTT observations of the GRB field, made two weeks after the GRB by the team led by the late Jan Van Paradijs, that included all the Italian *BeppoSAX* group, identified the host galaxy of the GRB, later confirmed with HST observations, which pointed for the first time in a direct and unambiguous way to the extragalactic origin of GRBs.

Two years later, the Italian team, among others (Stanek et al. 1999; Beuermann et al. 1999), had a relevant role in inferring the presence of relativistic jets in GRBs from the steepening of the optical light curve of the afterglow of GRB990510, observed with the ESO NTT, VLT and 3.6m telescope (Israel et al. 1999). Consistent with jetted emission, significant optical linear polarization at the 1.6-1.7% level was independently detected in the same GRB afterglow by two teams, one of which was under Italian leadership (Covino et al. 1999; Wijers et al. 1999).

GRB990123 confirmed a prediction that GRBs, besides exhibiting multi-wavelength afterglows, have prompt *simultaneous* very bright optical counterparts, called “optical flashes”, likely produced by internal or by reverse shocks, while afterglows are rather due to the interaction of the GRB blast-wave with the circumstellar medium. These optical flashes can be extremely bright (in GRB990123, the first one detected, it peaked at $V \sim 9$, Akerlof et al. 1999), but they are also dramatically ephemeral, so that only small, flexible robotic telescopes can observe and detect them. The record holder is GRB080913B, also called “naked-eye GRB”, because its optical flash reached the 5th magnitude. Its brightness made it suitable for very fast photometry with the Italian telescope REM at La Silla, equipped with the TORTORA instrument, that detected extremely rapid (time-scale of few seconds) optical variations correlated with the gamma-rays variations (Racusin et al. 2008). The bright optical afterglow was also the target of VLT UVES observations in Rapid Response

Mode, that started at 8.5 minutes after the GRB onset, when the magnitude was $R \sim 12$. In the UVES spectrum were detected absorption features belonging to the main system at $z = 0.937$ and divided in at least six components spanning a total velocity range of 100 km s^{-1} . These are estimated to be at distances between 2 and 6 kpc from the GRB site. Strong and variable Fe II fine structure lines were also observed, whose optical depth variation correlated with the UV/optical flux decrease, thus proving that the excitation of the observed fine structure lines is due to pumping by the GRB UV photons (D’Elia et al. 2009).

The host galaxies of GRBs bear the signatures of the stellar populations in which the explosion has taken place, and may therefore provide a way to characterize the class of progenitors of these powerful events, through the study of their basic properties, like luminosity, stellar mass, intrinsic absorption, star-formation rate, metallicity. The Italian astronomers have been responsible for many studies of the host galaxies of GRBs with ESO telescopes. Recently, a pilot study of GRB hosts with the newly deployed spectrograph X-Shooter, in the framework of the Italian-French Guaranteed Time program, underlined the potential of this new instrument in the accurate study of the metallicity of GRB hosts. The numerous lines that X-Shooter can detect make possible the application of a number of metallicity indicators and extend the investigation to the “redshift desert”, i.e. the interval $1 < z < 2$, for which the access to the optical range alone is not completely effective for host galaxy emission line detection (Piranomonte et al. 2011).

4. Gamma-Ray Bursts meet supernovae

One year after the outbreak of the GRB-afterglow era, an energetic Type Ic supernova³,

³ i.e., a core-collapse supernova whose progenitor has lost both its hydrogen and helium envelopes, to be distinguished from Type Ib supernova, with helium but no hydrogen and from Type II supernova, where both hydrogen and helium are kept before explosion.




Fig. 1. (left) Bolometric light curves of GRB and XRF supernovae at $z < 0.3$ (all observed with the ESO NTT and VLT) and of the “normal” (i.e., not accompanied by a GRB) Type Ic SN 1994I in rest frame (adapted from Melandri et al. 2012). For GRB/XRF supernovae, the explosion time is assumed to coincide with the GRB/XRF trigger time. The explosion date of SN 1994I was obtained from light-curve models. The supernovae associated with GRBs (GRB980425/SN 1998bw, 35 Mpc; GRB030329/SN 2003dh, $z = 0.168$; GRB021203/SN 2003lw, $z = 0.106$; GRB120422A/SN 2012bz, $z = 0.283$) are the most luminous (synthesized radioactive ^{56}Ni of 0.3–0.4 M_{\odot}) and are similar. The XRF supernovae (XRF060218/SN 2006aj, $z = 0.033$ and XRF100316D/SN 2010bh, $z = 0.059$) are less luminous. SN 2010bh is comparable in maximum luminosity to SN 1994I (10 Mpc), but faster in its rise to peak. This rapid rise time may indicate a strong asymmetry in the explosion. (right) Time evolution of the photospheric expansion velocities of GRB and XRF supernovae evaluated from the width of the SiII 6355Å spectral absorption line at various epochs (color coding as in left panel). The large velocities, especially at early epochs, imply high kinetic energies ($\geq 10^{52}$ erg). Note that SN 2010bh is the only supernova for which the width velocities were measured from the spectra and not evaluated with the aid of a model. The difference between the two methods amounts to about 20%, and the main uncertainty of the former method is related to the possible blending of the SiII 6355Å line with other transitions.

SN 1998bw, at 35 Mpc was detected and identified in the error box of the GRB980425, that had been detected by the *CGRO* BATSE and *BeppoSAX* GRBM and localized to arcminute precision by the *BeppoSAX* Wide Field Cameras (Soffitta et al. 1998; Galama et al 1998; Pian et al. 2000). That no optical afterglow was detected for this GRB, with the classical power-law time decay that astronomers had become accustomed to identify during one year of afterglow studies, came to a surprise⁴.

SN 1998bw, whose explosion time was consistent with the GRB trigger time within

⁴ GRB980425 exhibited no X-ray afterglow either, but its X-ray counterpart, which was identified with the X-ray emission from SN 1998bw, decreased by a factor of 2 in six months, and further decayed by a factor of 10 in the following 4 years (Pian et al. 2000; Kouveliotou et al. 2004).

~ 1 day, was exceptionally bright for a core-collapse supernova, having reached at maximum a luminosity comparable to that of a thermonuclear supernova (the synthesized radioactive ^{56}Ni , estimated from radiation transport models, is $\sim 0.4 M_{\odot}$), and even more importantly, it exhibited initial photospheric velocities of up to 30000 km/s, that implied a kinetic energy of 5×10^{52} erg, more than an order of magnitude larger than the typical 10^{51} erg (Mazzali et al. 2006a).

Later on, supernova features have been identified up to $z \sim 1$ in a large fraction of optical afterglows of long GRBs⁵ with well moni-

⁵ The majority of GRBs have durations longer than ~ 2 s (Kouveliotou et al. 1993) and are thus defined as long GRBs, as opposed to short or sub-second GRBs. Supernovae have been so far detected only for long GRBs.

tored light curves and accurately measured optical spectra as observed with ESO telescopes by European teams, including or led by Italian scientists (e.g., Hjorth et al. 2003; Della Valle et al. 2003; Malesani et al. 2004; Ferrero et al. 2006; Pian et al. 2006; Sparre et al. 2011; Bufano et al. 2012; Melandri et al. 2012). The GRB-supernova connection is best studied at $z \lesssim 0.3$, because the low distance allows spectroscopic monitoring of good signal-to-noise ratio and a full characterization of the supernova (see Figure 1). These supernovae are all of Type Ic, i.e. their spectra lack hydrogen and helium lines, and are therefore the explosion of highly stripped massive stars, that have entered the Wolf-Rayet phase.

Two of the GRBs associated with low-redshift well monitored supernovae are actually X-ray Flashes (XRFs), i.e. GRBs with softer spectra (i.e. peaking in the X-rays rather than the gamma-rays). Not only the synthesized ^{56}Ni and kinetic energies of GRB-supernovae are larger than in normal core-collapse supernovae⁶, but also their ejecta masses (Mazzali et al. 2006a). This points to progenitor main sequence masses of at least $20 M_{\odot}$ and, for the most massive GRB-supernovae, $40 M_{\odot}$ (Mazzali et al. 2006a;b; Bufano et al. 2012). The latter is considered to be the upper limit for the main sequence mass of a star undergoing supernova. Beyond this mass, the core should collapse directly to a black hole (e.g. Heger et al. 2003), perhaps after going through a brief (few seconds?) neutron star phase. In this case, if a GRB is formed, it will not be accompanied by a supernova, which may explain the lack of supernova detection, down to very deep photometric limits obtained with the VLT and HST, in the $z \sim 0.1$ GRB060614 and GRB060505 (Della Valle et al. 2006; Fynbo et al. 2006; Gal-Yam et al. 2006).

Finally, while GRB and XRF supernovae have maximum brightnesses in a range of 1.5 magnitudes (Figure 1), the energy outputs of their high energy events span four orders of

magnitude (Melandri et al. 2012, and references therein). This range is even wider if we include the higher redshift GRBs, whose promptly emitted gamma-ray energy is 10^{52} to 10^{54} erg. The wide interval of GRB gamma- and X-ray luminosities and the fact that local GRBs/XRFs are under-energetic point to the role of the viewing angle, that influences dramatically the jet emission via special relativity effects. While SN luminosities are much less affected by these, the explosion may be highly anisotropic, which will be reflected in the SN light curves and spectra, especially at late epochs ($\gtrsim 200$ days after explosion), when the inner ejecta are exposed and nebular, forbidden emission lines are formed. The profiles of these nebular lines, markedly those of oxygen and iron, have been studied with VLT FORS2 spectroscopy and they reveal details on the geometry of the explosion, on the degree of asymmetry and on the viewing angle (e.g. Mazzali et al. 2007b; Valenti et al. 2012).

The low redshifts at which GRB and XRF supernovae are best studied make also their close environments better targets for accurate studies. This is efficiently accomplished through space-resolved spectroscopy of a small region centered on the GRB site. An example is the study of the $z = 0.49$ host galaxy of GRB091127, associated with SN2009nz (Cobb et al. 2010; Berger et al. 2011), for which X-Shooter Integral Field Unit spectroscopy evidenced, via the construction of two-dimensional velocity, velocity dispersion, and star-formation rate maps, perturbed rotation kinematics and a star-formation rate enhancement consistent with the supernova position (Vergani et al. 2011).

5. Conclusions

Many supernova surveys currently active or coming on line will soon make the study of these explosions systematic and accurate and will at the same time guarantee large statistics (with many hundreds of supernovae detected per year) and high level of detail in the study of individual cases of particular and peculiar interest. These include the Palomar Transient Factory, the Panoramic

⁶ The properties of XRF-associated supernovae are intermediate between those of GRB-supernovae and normal Ic supernovae, see Figure 1.

Survey Telescope & Rapid Response System and the SkyMapper, all counting also on ESO telescopes for follow-up observations. Among these is the Public ESO Spectroscopic Survey of Transient Objects with the ESO NTT, led by S. Smartt and with a strong Italian participation. Similarly, the *Swift* satellite and other orbiting facilities for high energy astrophysics guarantee a rate of detection of GRBs and XRFs of about 100 events per year. A fraction of these can be followed up within few minutes by the VLT, thanks to the Rapid Response Mode, that will secure, after prompt identification via acquisition photometry, timely spectroscopy with the VLT UVES or X-Shooter. Later on, associated supernova search and study of the environment will take place, assuming adequate observing time can be devoted to these programs. Needless to say, besides clarifying existing problems, these observing programs will open new ones and will bring along the serendipitous detection of many transients of unexpected nature, that may create new interest and new research sub-branches in the two already very rich topics of supernovae and GRBs. This will also help shape up the characteristics and performances of the future extremely large telescopes and of their instruments.

Acknowledgements. I am grateful to A. Antonelli, S. Benetti, S. Covino, V. D'Elia, M. Della Valle, P. Ferrero, D. Malesani, N. Masetti, P. Mazzali, E. Palazzi and M. Turatto for their collaboration and for sharing with me their experiences and memories on GRB and supernova observational activity at ESO in the last 30 years. I would like to thank the organizer Vincenzo Mainieri for a very stimulating Symposium. In the last 10 years, the supernova and GRB research programs have benefited, among other sources of funding, from the dedicated European FP5 contracts for Research Training Networks HPRN-CT-2002-00303 (*The Physics of Type Ia Supernovae*) and HPRN-CT-2002-00294 (*Gamma-Ray Bursts: an Enigma and a Tool*), from INAF PRIN contracts in 2006, 2009, 2011, from PRIN MIUR 2005, and from ASI-INAF contracts I/023/05/0, I/088/06/0, I/016/07/0, I/009/10/0.

References

Akerlof, C., et al. 1999, *Nature*, 398, 400

- Benetti, S., et al. 2005, *ApJ*, 623, 1011
 Berger, E., et al. 2011, *ApJ*, 743, 204
 Beuermann, K., et al. 1999, *A&A*, 352, L26
 Bufano, F., et al. 2012, *ApJ*, 753, 67
 Cobb, B., et al. 2010, *ApJ*, 718, L150
 Costa, E., et al. 1997, *Nature*, 387, 783
 Covino, S., et al. 1999, *A&A*, 348, L1
 D'Elia, V., et al. 2009, *ApJ*, 694, 332
 Della Valle, M., et al. 2003, *A&A*, 406, L33
 Della Valle, M., et al. 2006, *Nature*, 444, 1050
 Ferrero, P., et al. 2006, *A&A*, 457, 857
 Fynbo, J., et al. 2006, *Nature*, 444, 1047
 Galama, T.J., et al. 1998, *Nature*, 395, 670
 Gal-Yam, A., et al. 2006, *Nature*, 444, 1053
 Hamuy, M., et al. 1996, *AJ*, 112, 2408
 Heger, A., et al. 2003, *ApJ*, 591, 288
 Hjorth, J., et al. 2003, *Nature*, 423, 847
 Israel, G., et al. 1999, *A&A*, 348, L5
 Kouveliotou, C., et al. 1993, *ApJ*, 413, L101
 Kouveliotou, C., et al. 2004, *ApJ*, 608, 872
 Lamb, D.Q. 1997, *PASP*, 107, 1152
 Maeda, K., et al. 2010, *Nature*, 466, 82
 Malesani, D., et al. 2004, *ApJ*, 609, L5
 Maund, J.R., et al. 2010, *ApJ*, 725, L167
 Mazzali, P.A., et al. 2006a, *ApJ*, 645, 1323
 Mazzali, P.A., et al. 2006b, *Nature*, 442, 1018
 Mazzali, P.A., et al. 2007a, *Science*, 315, 825
 Mazzali, P.A., et al. 2007b, *ApJ*, 661, 892
 Melandri, A., et al. 2012, *A&A*, in press
 (arXiv:1206.5532)
 Paczyński, B. 1997, *PASP*, 107, 1167
 Patat, F., et al. 1996, *MNRAS*, 278, 111
 Perlmutter, S., et al. 1998, *Nature*, 391, 51
 Perlmutter, S., et al. 1999, *ApJ*, 517, 565
 Phillips, M. M. 1993, *ApJ*, 413, L105
 Pian, E., et al. 2000, *ApJ*, 536, 778
 Pian, E., et al. 2006, *Nature*, 442, 1011
 Piranomonte, S., et al. 2011, *AN*, 332, 283
 Racusin, J.L., et al. 2008, *Nature*, 455, 183
 Riess, A., et al. 1998, *AJ*, 116, 1009
 Schmidt, B.P., et al. 1998, *ApJ*, 507, 46
 Soffitta, P., et al. 1998, *IAU Circ.* 6884
 Sparre, M., et al. 2011, *ApJ*, 735, L24
 Stanek, K., et al. 1999, *ApJ*, 522, L39
 Turatto, M., et al. 1996, *MNRAS*, 283, 1
 Van Paradijs, J., et al. 1997, *Nature*, 386, 686
 Valenti, S., et al. 2012, *ApJ*, 749, L28
 Vergani, S., et al. 2011, *A&A*, 535, A127
 Wijers, R.A.M.J., et al. 1999, *ApJ*, 523, L33