



Sloshing cold fronts in the IC1860 group

F. Gastaldello¹, L. Di Gesu^{1,2}, S. Ghizzardi¹, S. Giacintucci³, M. Girardi⁴,
E. Roediger⁵, and M. Rossetti^{3,1}

¹ IASF-Milano, INAF, via Bassini 15, Milan 20133, Italy
e-mail: gasta@lambrate.inaf.it

² Dipartimento di Fisica, Università degli Studi di Milano, via Celoria 16, 20133, Milan, Italy

³ Department of Astronomy, University of Maryland, College Park, MD 20742-2421, USA

⁴ Dipartimento di Fisica, Università degli Studi di Trieste, Sezione di Astronomia, via Tiepolo 11, 3

⁵ Hamburger Sternwarte, Universitaet Hamburg, Gojenbergsweg 112, D-21029 Hamburg, Germany

Abstract. We present a combined X-ray, optical, and radio analysis of the galaxy group IC 1860 using the currently available Chandra and XMM data, literature multi-object spectroscopy data and GMRT data. The Chandra and XMM data reveal two surface brightness discontinuities at 45 and 76 kpc shown to be consistent with a pair of cold fronts. We find evidence of a characteristic spiral pattern in the X-ray gas distribution, tilted with respect to the plane of the sky, indicative of a mainly line-of-sight off-axis merger responsible for the sloshing. We identify the likely perturber as an optically disturbed spiral galaxy, IC 1859. The peculiar velocity of the BCG with respect to the mean group velocity is another signature consistent with the sloshing scenario. The GMRT observation at 325 MHz shows extended radio emission: one component is contained within the inner cold front as seen in some galaxy clusters, whereas the second component is detached and in the vicinity of the spiral feature. The limits obtained from the NVSS at 1.4 GHz point to a very steep spectral emission coming from aged radio plasma. The evidence presented in this study is among the strongest supporting the currently favored model of cold front formation in relaxed objects and establishes the group scale as a chief environment to study this phenomenon.

Key words. clusters: general — X-rays: galaxies: clusters — hydrodynamics — galaxies: kinematics and dynamics — galaxies: clusters: individual (IC 1860 group)

1. Introduction

One of the most interesting discoveries by Chandra and XMM are the "cold fronts", sharp surface brightness discontinuities, interpreted as contact edges between regions of

gas with different entropies (see the review by Markevitch & Vikhlinin 2007). Cold fronts appear to be ubiquitous in galaxy clusters and they are found both in merging clusters and in cool core relaxed clusters. In relaxed clusters cold fronts are most likely induced by minor mergers that produce a disturbance on the gas in the core of the main cluster, displace

Send offprint requests to: F. Gastaldello




Fig. 1. *Left panel:* Exposure corrected mosaic of the MOS1 and MOS2 images in the 0.5-2 keV band smoothed on a $3''$ scale. *Right panel:* Map of the residuals between the XMM image and the best fit beta model for the data. The center of IC 1860 is shown by the black cross and the position and extent of the surface brightness edges by the red arcs. The “mushroom head” discussed in the text is the central region with white colour.

it from the center of the potential well, and decouple it from the underlying dark matter halo through ram pressure. The oscillation of the gas of the core around the minimum of the potential generates a succession of radially propagating cold fronts, appearing as concentric edges in the surface brightness distribution of the cluster. These fronts may form a spiral structure when the sloshing direction is near the plane of the sky and the merger has a non-zero angular momentum. When the sloshing direction is not in the plane of the sky concentric arcs are observed. The sequence of events is described in great detail in the simulations presented in Ascasibar & Markevitch (2006, AM06 hereafter). Another observable effect is the presence of a peculiar velocity of the central galaxy, if it sits at the cusp of the dark matter (DM) distribution and therefore starts oscillating with the DM peak after each subcluster flyby (AM06).

A number of relaxed, cool core clusters are hosts to faint, diffuse radio emission with a radius comparable to the size of the cool core and with a steep spectrum. These sources are called mini radio-halos and they are relatively rare, with only few clusters with confirmed detections (see Ferrari et al. 2008, for a review). The sloshing motions can produce turbulence in the cluster core. Mazzotta & Giacintucci (2008) discovered spatial correlations between

mini radio-halos emission and cold fronts in the X-ray images of two clusters, with the mini-halos contained within the region confined by the cold fronts and they suggested that the correlation arises from the turbulence generated by the sloshing motions. ZuHone et al. (2013) tested the hypothesis with high resolution MHD simulations of gas sloshing confirming that re-acceleration of relativistic electron seeds (e.g., from past AGN activity) can produce radio mini-halo emission within the envelope of the sloshing cold fronts.

Sloshing cold fronts have been detected also at the smaller mass scales of groups of galaxies (e.g. Gastaldello et al. 2009, G09 hereafter). At this mass scale even a single massive galaxy can be the responsible perturber and the X-ray signature of sloshing cold front is a signature of the gravitational interaction between the galaxy (and its DM halo) and its host group. Here we present the results of the available Chandra and XMM data, optical and radio GMRT data for the IC 1860 group, a nearby group of galaxies at $z=0.022$.

2. XMM and Chandra observations of IC 1860

The XMM image in Fig.1 (left panel) reveals the presence of a brightness discontinuity in the NW and a surface brightness extension




Fig. 2. XMM surface brightness profiles in two sectors, one including the outer cold front and one shown for comparison. Dots correspond to the sector PA 340°-70° and diamonds to the comparison sector 70°-160°. PA are measured from the W direction.




Fig. 3. XMM temperature profiles along the sectors containing the cold fronts. The vertical dashed lines mark the position of the inner and outer cold fronts. Circles refer to the 220°-250° sector hosting the inner cold front, diamonds to the 340°-70° sector hosting the outer cold front.

from the core to the SE which also ends in a surface brightness discontinuity. A more detailed view is provided by the surface brightness residual map between the XMM data and the best fit one-dimensional β -model in the right panel of Fig. 1. A characteristic spiral pattern is seen as a surface brightness excess even though the qualitative impression is that the overall morphology is tilted with respect to the plane of the sky, with the head of the mushroom structure seen in many hydrodynamic simulations (AM06) pointing towards the observer. The surface brightness profiles of the Chandra and XMM data extracted from selected angular sectors mark the position of the surface brightness discontinuities: at a radius of 170'' (71 kpc) for the outer edge in the NW and at 100'' (45 kpc) for the inner edge in the SE. In Fig. 2 we show the XMM data highlighting the outer cold front. In Fig. 3 we present the results of the XMM spectral fits (Chandra gives consistent results) in the radial bins inside and outside the edges. For the inner edge a clear temperature increase across the edge is detected, as expected for a cold front; there is no statistically significant change in tempera-

ture across the outer edge. However taking into account projection effects the picture is consistent with cooler gas inside the edge; a similar behavior has been seen for the outer cold fronts in NGC 5044 (G09) and NGC 5846 (Machacek et al. 2011, M11 hereafter). The emission in the SE leading to the inner cold front is cooler than the surroundings.

3. Radio and optical data

The GMRT radio image at 325 MHz (radio contours shown in the left panel of Fig. 4) reveals an extended feature which originates at the IC1860 center and extends from the center towards SE. This radio emission is confined within the inner cold front and traces the "mushroom head" feature interpreted as the tip of the slushing spiral. A second patch of faint radio emission is visible 2 arcmin SW of the central source. Neither of these extended features are visible in the 1.4 GHz VLA image by Dunn et al. (2010), indicating very steep spectral indexes for these features. The radio emission comes probably from aged radio plasma, perhaps associated with a past episode of ac-




Fig. 4. *Left panel:* GMRT radio contours at 325 MHz overlaid on the map of X-ray residuals (right panel of Fig.1). The restoring beam is $30'' \times 29''$, in p.a. 6° (white ellipse). The rms noise level is $0.6 \text{ mJy beam}^{-1}$. Contours are spaced by a factor of 2 starting from 2 mJy beam^{-1} . The emission which can be seen in the South is coincident with an X-ray point source. *Right panel:* DSS image of the central region of the IC 1860 group.

tivity of the central radio galaxy. In the optical band we collected 74 member galaxies with known velocity within 1.6 Mpc of IC 1860. The mean group velocity is $6753 \pm 44 \text{ km/s}$ with a dispersion velocity of $401 \pm 46 \text{ km/s}$. The peculiar velocity of the central galaxy IC 1860 with respect to the group mean is 113 km/s significant at the 2.3σ level. We identified as the possible perturber IC 1859, a Seyfert 2 galaxy with a peculiar optical morphology showing a pair of spurs in the eastern spiral arm (right panel of Fig.4). It is at a projected distance of 176 kpc from IC 1860 and it has a velocity difference of $-880 \pm 67 \text{ km/s}$ with respect to IC 1860 and of $-767 \pm 78 \text{ km/s}$ with respect to the group mean velocity.

4. Conclusions

The analogy of the IC 1860 group with other groups, like NGC 5044 (G09) and the NGC 5846 group with the rich, barred spiral NGC 5850 identified as the perturber (M11), is tantalizing. The case study presented in this paper shows the rich phenomenology and the

constraining power that X-ray, optical and radio data can provide in assessing the dynamical state and past merging history of galaxy groups and their galaxies. We have provided the strongest evidence to date that cold fronts in groups exist and have similar properties to those in massive clusters. This is critical confirmation of the currently favored model of cold fronts formation in relaxed objects.

References

- Ascasibar, Y. & Markevitch, M. 2006, ApJ, 650, 102
- Dunn, R. J. H., et al. 2010, MNRAS, 404, 180
- Ferrari, C., et al. 2008, Space Sci. Rev., 134, 93
- Gastaldello, F., et al. 2009, ApJ, 693, 43
- Machacek, M. E., et al. 2011, ApJ, 743, 15
- Markevitch, M. & Vikhlinin, A. 2007, Phys. Rep., 443, 1
- Mazzotta, P. & Giacintucci, S. 2008, ApJ, 675, L9
- ZuHone, J. A., Markevitch, M., Brunetti, G., & Giacintucci, S. 2013, ApJ, 762, 78