



# A Spanish-VO tool to estimate the spatial distribution of galactic interstellar extinction

C. Morales-Durán<sup>1</sup>, C. Rodrigo-Blanco<sup>1</sup>, P. Cabo-Cubeiro<sup>2</sup>,  
F. Llorente-de Andrés<sup>1</sup>, and E. Solano-Márquez<sup>1</sup>

<sup>1</sup> Centro de Astrobiología, CSIC-INTA, ESAC campus, apartado postal 78, 28 691 Villanueva de la Cañada, Madrid, Spain

<sup>2</sup> INTA, apartado postal 78, 28 691 Villanueva de la Cañada, Madrid, Spain e-mail: morales@cab.inta-csic.es

**Abstract.** The interstellar medium in our Galaxy is very inhomogeneous. The interstellar gas and dust are distributed irregularly in clouds, filaments arcs, bubbles and superstructures of different density from very high to extremely low. The knowledge of the spatial distribution of these features is of capital importance for different areas of the galactic and extragalactic astronomy. We present here a tool, developed in the frame of the Spanish Virtual Observatory, to know the variation of the extinction in any observing direction.

**Key words.** Stars: distances, extinction – ISM: extinction – Galaxy: local interstellar matter – Galaxy: open clusters and associations

## 1. Introduction

Maps of the spacial distribution of interstellar reddening traditionally were based in individual stellar data (FitzGerald,1968; Neckel et al.,1980; Arenou et al, 1990) but, due to the scarcity of good quality distance measurements, many theoretical models have been developed which must be tested with real



**Fig. 1.** Entering object query or coordinates in the SVO tool “Extinction Map”.  
data. The huge quantity of observational data

(Hipparcos, Tycho, SDSS, ASCC,...), from ground and from the space, published in the last few years will make now possible this effort. We are going to make all this available data accesible from the spanish virtual observatory through a page with easy and quick access. We present here a tool to visualize the distribution of the interstellar extinction. Using published photometric, spectroscopic and trigonometric data we have built a database of published and calculated by us distances and reddenings of 200 000 stars and growing. Open and globular clusters are added, being part of the data base. In the framework of the Spanish Virtual Observatory we have developed a tool called “Extinction Map” that allows queries to the database by name or coordinates and provides as output, extinction-related data (distance,  $A_v$ ,  $R_v$  and  $E(B-V)$  . If

Extinction map  
(Morales-Delgado et al., 2009)

Object	Position		Distance		Search	Extinction	(?)
	RA : J2000	Dec : J2000	RA : J2000	Dec : J2000	Dist (kpc)	E(B-V)	Plot (°) Area

54 files found.

Name	cat	RA	Dec	$\theta$	Dist	$V_{mag}$	E-V	E(B-V)	Rv	A <sub>v</sub>	SFR
HD15752	8	383250	582420	001126	1.932	8.74	0.49			2.42	30100
BD+57588	8	382119	583534	001941	2.999	10.11	0.61			2.7	20110
HD15440	11	375345	585129	002003	3.229	8.076	1.42	0.26		0.84	62.H
HD15620	8	381311	575545	002020	2.292	8.35	0.92			3.01	11180
HD15620	1	381313	575545	002020	2.09	8.35	0.92	0.95	2.867158	2.72	B8Iab
HD15630	11	381313	575545	002020	1.929	8.25	0.82	0.95		3.12	B9Iab
HD15619	11	381243	583635	002050	0.328	8.076	1.039	0.069		0.24	G9.H
HD16025	8	391144	580713	002938	0.341	8.97	0.17			0.77	S0I90
HD16025	11	391146	580712	002938	0.421	8.97	0.17		0.27	0.86	B9.V
HD16025	1	391146	580712	002938	0.4	8.97	0.17	0.245	2.849715	0.698	B9.V
US20502	8	372135	581254	002950	2.857	9.86	0.48			2.32	30110
BD+57593	8	383616	574820	003452	2.4	9.82	0.61			2.73	30110
BD+57593	11	385800	574721	003534	1.955	9.92	0.61	0.87		2.8	B1.H
VV4299e	1	375820	574151	003538	1.9	9.98	0.79	0.85	2.831858	2.4	B6Ia
HD15497	8	375618	574150	003538	1.967	7.03	0.762			2.61	10190
6.10092134	8	383822	585034	003612	2.4					2.43	99990
HD15316	8	372940	574914	003658	1.851	7.23	0.77			2.36	11230
HD15316	11	372938	574914	003658	1.848	7.227	0.77	0.71		2.35	A3.Iab
HD15964	11	390323	574341	004014	0.489	8.88	0.09	0.2		0.64	B9.V
HD16012	8	390852	574344	004222	0.481	8.48	0.07			0.66	30190
HD16012	11	390853	574344	004224	0.664	8.42	0.075	0.175		0.98	B9.H
BD+57579	1	373330	574030	004234	1.54	10.09	0.49	0.755	2.768854	2.08	B1.V
BD+57579	11	373330	574030	004234	1.374	10.09	0.49	0.75		2.41	B1.V
BD+57579	8	373304	573950	004318	1.968	10.09	0.49			2.59	S0I10
HD15690	8	382326	573236	004325	1.564	9.82	0.57			2.63	S0I10
HD15690	2	382311	573214	004347	2.488	8	0.66	0.83	2.92	2.42	B2b
HD15690	1	382311	573214	004347	1.64	8.01	0.67	0.85	2.765076	2.35	B2b
HD15690	8	382310	573213	004348	1.539	8.005	0.658			2.63	12115
BD+09057	8	381810	573015	004541	2.25					2.4	B90
HD23671	8	385426	572913	005024	1.799	9.55	0.5			2.4	40310
HD23671	1	385427	572913	005025	1.56	9.55	0.5	0.785	2.826144	2.23	B1.V

**Fig. 2.** Partial list of the objects found by the SVO tool “Extinction Map” around the entered coordinates or object name.

the queried object is not in the database, the system returns the information of all the stars located within a cone of variable radius.

## 2. Query by coordinates

We show here a preliminary example of a query. You enter the coordinates of the region of interest (Fig. 1) and the SVO tool provides you with a list of the stars present in the database inside a cone of radius chosen (Fig. 2). Also it provides with a representation of visual absorption versus distance and a second representation of the stellar coordinates (Fig. 3).

You may find in your search that certain stars have been studied by different researchers at different times or with different


methods. If uncertainties are provided you can delete from the list the worst values and made the representation again, but uncertainties are only provided for the very recent observations. Clicking on the column cat you may find the origin of the data and information about the original values. In the figure (Fig. 4) we have the Aitoff projection of the stars present by today on the database.

Many more published catalogues and our own computed distances and reddenings will be soon implemented and will be continuously implemented.


## 3. Potentiality

The interstellar extinction law shows a sharp rise in the extinction value from the visible towards the ultraviolet. Being able to know the proper amount of reddening in front of any astronomical object observed in the optical and specially in the UV is fundamental and more and more important as we go towards smaller wavelengths. Other potential users of this tool are astronomers whose objects by its intrinsic characteristics are not suitable to obtain easily the E(B-V) and therefore its distance. A preliminary version of this tool is available as a web page (<http://svo2.cab.inta-csic.es/theory/exmap/>) and also as a ConeSearch Virtual Observatory service.

*Acknowledgements.* This research has made use of the Spanish Virtual Observatory supported from the Spanish MINECO through grant AyA2011-24052; F. Llorente-de Andrés acknowledges support from the ESAC Faculty.



**Fig. 3.** Left: Representation of visual absorption versus distance, and Right: representation of the stellar coordinates.



**Fig. 4.** Aitoff projection of the stars present by today on the database.