



CO concentration in the upper stratosphere and mesosphere of Titan: non-LTE analysis of VIMS dayside limb observations at $4.7 \mu\text{m}$

F. Fabiano^{1,2}, M. Lopez-Puertas³, A. Adriani⁴, M.L. Moriconi⁴, E. D'Aversa⁴,
B. Funke³, M.A. Lopez-Valverde³, M. Ridolfi¹, and B.M. Dinelli²

¹ Dept. of Physics and Astronomy, University of Bologna, Viale Berti Pichat 6/2, 40127 Bologna, Italy

² CNR Institute of Atmospheric Sciences and Climate (ISAC-CNR), Via Piero Gobetti 101, 40129 Bologna, Italy

³ INAF–Institute for Space Astrophysics and Planetology, Via Fosso del Cavaliere 100, 00133 Roma, Italy

⁴ Instituto de Astrofísica de Andalucía CSIC, Glorieta de la Astrónoma, s/n, E-18008, Granada, Spain

Abstract.

During the last 20 years, many works have focused on the atmospheric concentration of CO on Titan, giving contradictory results. In particular, no measurement of the CO abundance above 300 km has been done yet, due to the faint emission of CO above that altitude. On the other hand, such a study is particularly awaited as a confirmation of photochemical models that predict a uniform volume mixing ratio of CO in the whole Titan's atmosphere. Moreover, given that CO is the main reservoir of oxygen in Titan's atmosphere and its presence is linked to water, the matter is of astrobiological interest too. The analysis of VIMS (Visual & Infrared Mapping Spectrometer, onboard Cassini) daytime limb measurements of Titan at $4.7 \mu\text{m}$, corresponding to the vibrational bands of CO, allows such a study: CO molecules are significantly excited by solar radiation and the otherwise faint infrared signal of the upper atmosphere is large, allowing to probe this region too. On the other hand, the strong non-LTE behavior of CO infrared emission above 200 km, strongly coupled to N₂ and CH₄ vibrational levels, represents an extra complexity of the data analysis. In order to deal with CO non-LTE emission, we have developed a non-LTE excitation - de- excitation model for the first two CO vibrational levels of the two most abundant isotopes. We consider the contribution of absorption of radiation in the fundamental, first hot and first overtone bands and evaluate the possible contribution of different collisional processes, mainly the coupling with the first excited state of N₂ and with several levels of CH₄. The solution of the non-LTE problem is obtained (Funke et al. 2012) through GRANADA, a non-LTE population algorithm based on Lambda iteration resolution strategy, eventually coupled to a Curtis-Matrix type approach. The result shows a significant over-population of the first

vibrational level above 300km and of the second level in the whole atmosphere. The possibility of further pathways for CO(1) (de)-excitation, mainly through vibrational energy transfer from - to the excited states of CH₄, is explored and seems to be crucial for establishing its population. We then analyze the spectra acquired by VIMS between 2004 and 2013 in the 4.7 μm region, for daytime conditions. Considering the non-LTE populations obtained above, the retrieval of CO concentration is performed with the aid of Geofit Broad Band, a non- LTE line-by-line radiative transfer code coupled to a bayesian inversion method, initially developed for the Earth's atmosphere and later adapted to other planetary atmospheres (Adriani et al. 2011). The low signal-to-noise ratio and spectral resolution of the instrument considerably complicate the data analysis. Moreover, the scattered solar radiation is not negligible below 350 km and dominates over atmospheric emission below 200 km, and hence it is taken into account as well. CO relative abundance profiles are finally obtained and discussed in the light of photochemical models predictions.

Acknowledgements. Special thanks are given to the Cassini/VIMS team for planning the observations. The ISAC-CNR and IAPS-INAF teams have been funded by the Cassini project of the Italian Space Agency under the contract ASI I/015/09.

References

- Adriani, A., Dinelli, B.M., López-Puertas, M., et al. 2011, *Icarus*, 214, 584
Funke, B., López-Puertas, M., García-Comas, M., et al. 2012, *J. Quant. Spec. Radiat. Transf.*, 113, 177