

High-amplitude δ Scuti stars in the Galactic Bulge from the OGLE-II and MACHO data

A. Pigulski, Z. Kołaczkowski, T. Ramza, and A. Narwid

Instytut Astronomiczny Uniwersytetu Wrocławskiego, Kopernika 11, 51-622 Wrocław, Poland, e-mail: pigulski@astro.uni.wroc.pl

Abstract. Searching for main-sequence pulsators, we analyzed photometry of $\sim 200,000$ variable star candidates from the OGLE-II Galactic fields, finding 193 high-amplitude δ Scuti stars. This doubles the number of known stars of this type. The MACHO data, available for half of stars, were also analyzed. In our sample of the HADS stars, we found 50 multiperiodic objects, including 39 that have period ratios in the range of 0.76–0.80, an indication of the radial fundamental and first-overtone pulsation. We discuss the resulting Petersen diagram for these stars in view of the period ratios predicted by models. Except for stars showing pulsations in the radial fundamental mode and first overtone, we find the evidence for higher radial overtones and non-radial modes in the analyzed sample of multiperiodic HADS stars.

Key words. Stars: δ Scuti, Stars: pulsations

1. Introduction

During the last several years, large photometric time-series data have been made available. Using massive photometries from the OGLE, MACHO, ASAS and NSVS projects, we started a systematic study of the Galactic main-sequence pulsators. This paper presents the results of the search for high-amplitude δ Scuti (hereafter HADS) stars in the catalogue of $\sim 200,000$ variable star candidates published by Woźniak et al. (2002). This catalogue contains I -filter photometry carried out between 1997 and 2000 within the OGLE-II project (Udalski et al. 1997). The observations covered about 11 square degrees in 49 Galactic fields. The OGLE-II data were

supplemented by the MACHO observations (Allsman & Axelrod 2001), if available.

HADS stars form a subgroup of δ Scuti stars with large amplitudes (exceeding ~ 0.3 mag in V band), non-sinusoidal light curves and small rotational velocities (see, e.g., McNamara 1997). There is, however, no clear separation between them and δ Scuti stars with small amplitudes. HADS stars pulsate mainly in radial modes, but there is a growing evidence that non-radial modes with small amplitudes can also be unstable in these stars. Population II HADS stars are called SX Phoenicis stars and are discovered mainly in the globular clusters. At present, we know about 150 HADS stars in the Galactic field (Rodríguez et al. 2000) and about 200 SX Phe stars in globular clusters and nearby galaxies

Send offprint requests to: A. Pigulski

(Rodríguez & López-González 2000, Clement et al. 2001).

The MACHO microlensing survey data in Galactic fields were already used to search for the HADS stars by Alcock et al. (2000), who found 90 stars of this type, including 18 multiperiodic objects.

2. Analysis

The first two steps in our analysis were done automatically. All stars from the catalogue of variable star candidates of Woźniak et al. (2002) were analyzed by means of the Fourier periodogram. Consecutive detection and subtraction of up to five periodic terms was performed. The Fourier spectra were calculated up to 40 d^{-1} . Next, the star that appeared to be variable was classified by means of the detected period(s), amplitude(s) and the results of Fourier decomposition. This information was used to select stars that were subsequently analyzed interactively. At this step, it was decided whether the star was included in the list of the HADS stars or not. We adopted the following definition of a HADS star: it is a star with period shorter than 0.25 d for which at least one harmonic of the main mode was detected and which is not an RR Lyrae or W UMa star. The latter two types were distinguished from the HADS stars by means of Fourier coefficients and/or visual inspection of the light curves. In total, 193 HADS stars were found.

It is interesting to mention that this procedure allowed us to discover also other main-sequence variables or at least candidates. Many of them are multiperiodic. We have found about 230 β Cephei and low-amplitude δ Scuti stars (Narwid et al., these proceedings), and about 600 candidates for SPB and γ Doradus stars.

As mentioned in the Introduction, for stars classified as HADS the photometry from MACHO database was retrieved, if present, and analyzed in a similar way. It turned out that about a half of HADS stars detected in the OGLE data have MACHO photometry, but only 15 of them were found by Alcock et al. (2000).




Fig. 1. The histogram of periods detected in 193 HADS stars found in the OGLE-II Galactic fields.




Fig. 2. The semi-amplitudes of the main components of the highest-amplitude mode in the MACHO blue filter, $A_1(V_M)$, plotted against semi-amplitudes of the component in the MACHO red filter, $A_1(R_M)$. The continuous line is the best-fit of the amplitude ratio equal to 1.45 ± 0.01 . The dashed line denotes $A_1(V_M) = A_1(R_M)$.

3. Results

The histogram of periods detected in all 193 HADS stars is shown in Fig. 1. We see that it peaks for periods equal to $0.08\text{--}0.1 \text{ d}$ and that no single period longer than 0.2 was found.




Fig. 3. Petersen diagram for double and multiple-mode HADS stars with period ratio in the range between 0.74 and 0.81. Filled circles: 25 stars listed by Poretti et al. (2005), open circles: the remaining seven stars discovered by Alcock et al. (2000), but not included by Poretti et al. (2005), crosses: 39 stars from this study, open triangles: six double-mode SX Phe stars from NGC 3201 and NGC 5466. Two vertical lines join points for the same star, but two possible values of P_1 .

On the other hand, the shortest period we detected equals to about 0.03 d. For about 90 stars common to OGLE-II and MACHO data we were able to find the ratios of amplitudes of the main modes. The mean ratios of amplitudes were equal to $A_1(V_M)/A_1(I) = 1.80 \pm 0.03$, $A_1(R_M)/A_1(I) = 1.25 \pm 0.02$, and $A_1(V_M)/A_1(R_M) = 1.45 \pm 0.01$, where A_1 stands for the amplitude of the main component of the highest-amplitude mode in the Fourier decomposition. The MACHO blue and red filters are denoted as V_M and R_M , respectively. The $A_1(V_M)$ vs. $A_1(R_M)$ dependence is shown in Fig. 2. The small r.m.s. errors of the ratios can be easily explained, since the modes detected in the HADS stars are mostly radial, and therefore their amplitudes should change with wavelength in a similar way.

In our sample of 193 HADS stars, 52 stars show multiperiodic behaviour. Of them, 40

stars are double-mode pulsators, while the remaining 12 have three or even more modes excited. Double-mode HADS stars are very useful, as if the period ratio falls in the range of 0.76–0.79, the modes can be interpreted as radial: fundamental and first overtone (Petersen & Christensen-Dalsgaard 1996).

Double-mode HADS stars were recently studied by means of the Petersen diagram by (Poretti et al. 2005), who provided an updated list of 25 known stars of this type. These stars are shown with filled circles in Fig. 3 which is the Petersen diagram ¹ showing all period ratios (with respect to the period of the main

¹ In principle, the Petersen diagram is used to plot the ratio of the period of the first overtone and fundamental radial modes, i.e. P_1/P_0 , versus P_0 . However, we use this diagram in a broader context, plotting the ratio of a period of a given mode and the period of the main mode, presumably the fundamen-

mode, P_0) in the range between 0.74 and 0.81. Seven double-mode HADS stars in the Poretti et al.'s (2005) list were found in the MACHO data. However, there are seven more stars, not included by these authors, that have period ratios not very close to 0.77, but still in the range covered in Fig. 3. These stars are shown with open circles in Fig. 3. In fact, there is one star, MACHO 120.21785.976, with the reported period ratio of 0.71199 (Alcock et al. 2000). If, however, the alias frequency of the f_2 mode is taken as the true one, the period ratio goes very close to the canonical value and amounts to 0.77141. We plot the latter value in Fig. 3.

We add 39 stars to this sample (crosses in Fig. 3), which increases the number of known double-mode HADS stars more than twofold. The new stars populate mostly the short-period region of the diagram, where no star of this type was found by Alcock et al. (2000), apparently due to some selection effects in their analysis.

In order to explain the Petersen diagram for the HADS stars, Poretti et al. (2005) used appropriate stellar models for different masses and metallicities to calculate theoretical values of P_1/P_0 . A similar work has been done earlier by Petersen & Christensen-Dalsgaard (1996). These calculations show that both mass and metallicity influences P_1/P_0 . The effect of mass is largest for periods longer than ~ 0.12 d where lower mass leads to the lower value of P_1/P_0 . On the other hand, the metallicity is most important in the short-period range. The smaller metallicity the larger P_1/P_0 . The metallicity effect is, in general, confirmed by the observations of double-mode SX Phe stars in globular clusters. This can be judged from Fig. 3 where, for comparison, we show (open triangles) period ratios for six double-mode SX Phe stars which are the members of two metal-poor Galactic globular clusters: NGC 3201 (Mazur et al. 2003) and NGC 5466 (Jeon et al. 2004).

tal radial one. This does not mean that in each case we deal with two radial modes. In fact, for some stars, we use the Petersen diagram to claim the opposite (see text).

Consequently, we may conclude that some double-mode HADS stars we found in the OGLE-II data could be SX Phe stars, especially those that have periods shorter than 0.1 d and $P_1/P_0 > 0.775$. It is also obvious that some of the ratios shown in Fig. 3 fall outside the range of P_1/P_0 allowed by models. There are at least two plausible explanations of this fact. For ratios larger than 0.78 we may deal with a pair of consecutive radial overtones (see Templeton et al. 2002), radial and non-radial mode or even two non-radial modes. We see that for at least two stars shown in Fig. 3, this must be true, as all their periods cannot be explained in terms of only radial modes.

In addition to the modes shown in Fig. 3, we found in multiperiodic HADS stars modes that are good candidates for higher radial overtones and non-radial modes. There is also an evidence for period changes in some HADS stars we found. This conclusion comes from the fact that for these stars a peak at frequency very close to the main mode appears in the spectrum of residuals.

Acknowledgements. This work has been supported by the KBN grant No. 1 P03D 016 27.

References

- Alcock, C. et al. 2000, ApJ, 536, 798
 Allsman, R.A. & Axelrod, T.S. 2001, astro-ph/0108444
 Clement, C.M. et al. 2001, AJ, 107, 2131
 Jeon, Y.-B. et al. 2004, AJ, 128, 287
 Mazur, B., Krzemiński, W. & Thompson, I. 2003, MNRAS, 340, 1205
 McNamara, D.H. 1997, PASP, 109, 1221
 Petersen, J.O. & Christensen-Dalsgaard, J. 1996, A&A, 312, 463
 Poretti, E. et al. 2005, A&A, 440, 1097
 Rodríguez, E., & López-González, M.J. 2000, A&A, 359, 597
 Rodríguez, E., López-González, M.J. & López de Coca, P. 2000, A&AS, 144, 469
 Templeton, M., Basu, S., & Demarque, P. 2002, ApJ, 576, 963
 Udalski, A. et al. 1997, AcA, 47, 319
 Woźniak, P.R. et al. 2002, AcA, 52, 129